

407786

REPOSITORY Brookhaven National LAB  
COLLECTION MEDICAL RECORDS Dept  
BOX No. NA  
FOLDER NA

PRELIMINARY REPORT

RONGELAP REASSESSMENT PROJECT

APRIL 20, 1988

TO: The President and Congress of the United States  
FROM: Henry I. Kohn, Referee

Rongelap Reassessment Project  
1203 Shattuck Av., Berkeley, CA 94709  
(415) 526-0141

The Medical Research Center  
Brookhaven National Laboratory  
Upton, L. I., New York

## ABSTRACT

This preliminary report provides the basis for testimony to be given on April 26, 1988, before the House Appropriation Committee on Interior, Representative Sidney Yates, Chairman.

It was considered important for both the Congress and the Rongelap people to present an overview of the material now available rather than to wait until all questions have been answered. Meeting the hearing date has involved some last minute pressures. The final report will probably be issued within 2 - 3 months.

The chief conclusion is that, based on the estimation of adult dosage, Rongelap Island may be resettled now. That conclusion, however, presupposes certain conditions for living which are set out and discussed in Section 5 (which may be read without reference to the rest of the Report).

The chief unsettled point is the dose to infants; it is currently under review.

Another unsettled point is the transuranic dosage (plutonium-293, -240, americium-241).

It is important to bear in mind that the dosage under discussion is that from continued residence on Rongelap Island from 1978 (or the present), onwards. This adult dosage over the next 30 years is estimated to be no more than 1 to 2% of that experienced from fallout in 1954 from the Bravo shot. The historical data included in the Report are of interest for general orientation.

As referee, I am solely responsible for the contents of this report. However, two consultants have strongly objected to major portions of it and I am therefore putting their comments together, in their entirety, in Note 13. For comparison, I suggest that they be read in conjunction with Section 5 of the Report (Discussion and Recommendations).

## TABLE OF CONTENTS

ABSTRACT	1
1. INTRODUCTION	3
1.1 Task	3
1.2 Procedure	3
2. BACKGROUND -- THE RONGELAP EXPERIENCE	7
2.1 Bravo test -- 1954	7
2.2 Return to Rongelap -- 1957	10
2.3 Rongelap 1957 -- 1987	10
3. REASSESSMENT	15
4. DOSE	19
4.1 External Dose	20
4.2 Internal Dose: Lawrence Livermore	23
4.3 Internal Dose: Brookhaven	29
4.4 Infant Dosage	32
4.5 Dose Summary	33
5. DISCUSSION AND RECOMMENDATIONS	36
5.1 Assumptions	36
5.2 Infant Dosage	37
5.3 Plutonium	37
5.4 Monitoring and Health Programs	37
5.5 Rehabilitation of Soil	38
6. REFERENCES	41
7. NOTES	46

## 1. INTRODUCTION

### 1.1 Task

Rongelap Atoll was contaminated with radioactive fallout in 1954 as a result of the Bravo thermonuclear test-shot at Bikini, 130 miles away. In 1978, to inform the Rongelap people of the extent of residual contamination 24 years later and of its potential effects upon their health, DOE (Department of Energy) surveyed the region and subsequently issued a specially prepared book report in Marshallese.

The book was entitled, The Meaning of Radiation for Those Atolls in the Northern Part of the Marshall Islands that were Surveyed in 1978, and was published in 1982. (We shall refer to it as DOE-1982.) The first part dealt in general with radiation and fallout, and how they might affect plants, animals and man. The situation at Rongelap was dealt with specifically on pages 38 - 39. (Note 1)

DOE's assessment of Rongelap Island was not accepted by the Rongelap people, so much so that in 1985 the residents abandoned their homes and moved to Majiето in Kwajalein Atoll.

The U. S. Congress, therefore, provided for an independent assessment of DOE's conclusions for Rongelap Island in the Compact of Free Association Act of 1985 (U.S. Public Law 99-239, section 103(i); see Note 2). The functions of the present report are therefore as follows:

"[The referee shall] review the data collected by the Department of Energy relating to the radiation levels and other conditions on Rongelap Island resulting from the thermonuclear test...The purpose...shall be to establish whether the data cited in support of the conclusions as to habitability of Rongelap Island as set forth in the [book] ...are adequate and whether such conclusions are supported by the data....If...the data are inadequate to support...habitability...the government of the Marshall islands shall contract...[for]...a complete survey...[and for recommendations of]...the steps needed to restore habitability..."

### 1.2 Procedure

The DOE-1982 book now under review was discussed with its senior author, Dr. William Bair (Pacific Northwest Laboratories, Richland, Washington 99352), and Dr. Bair has read the parts of this Report referring to it. Dr. William Robison (Environmental Sciences Division, Lawrence Livermore Laboratory, Livermore CA 94550), who supplied the field data was also interviewed and has read this Report.

Relevant Rongelap studies that were supported by DOE at Brookhaven National Laboratory (Upton, New York 11973), were discussed with Dr. William H. Adams, (Medical Department) and Mr. E. Lessard (Safety & Environmental Protection Division). The citation of their work in this Report has been checked by them.

Additional information from DOE-supported laboratories that became available after DOE-1982 had been written was made available to us by Adams, Lessard and Robison. Also, we have taken a number of samples in the field and have had them analyzed independently.

Other sources of information in the international literature have been used and are cited in the text.

We have also discussed from time to time various matters relating to the Report, or the progress made in developing it, with the Rongelap people or their representatives, including Senator Jeton Anjain, P.O. Box 1006, Majuro, Republic of the Marshall Islands, 96960.

We have also consulted Mr. Peter Oliver, Special Assistant for Compact Affairs, Republic of the Marshall Islands, P.O. Box 15, Majuro, 96960.

The Reassessment Report (the present document) was written by Henry I. Kohn in his capacity as Referee under contract with RepMar. The opinions and statements made are therefore his responsibility. The task, however, was greatly facilitated by employing an international panel of experts, selected so as to represent a variety of overlapping specialties that would cover the problems under examination.

If they chose to do so, the consultants who were still in disagreement with the final draft of the Report (having discussed earlier versions with Dr. Kohn), were asked to write brief notes on their own views to be mentioned in the text and to be included as footnotes or among the "Notes to the Text". The absence of such comment, however, does not necessarily indicate agreement with the entire text. A major commentary by Dr. Bertell and Mr. Franke is given in Note 13.

The following scientists participated in the Project.

Referee

HENRY I. KOHN, Ph.D., M.D. (radiation biology) Gaiser Professor  
Emeritus of Radiation Biology, Harvard Medical School; Chairman,  
Bikini Atoll Rehabilitation Committee; 1203 Shattuck Ave., Berkeley  
CA 94709 (415-526-0141)

Consultants

S. J. ADELSTEIN, M.D., Ph.D. (nuclear medicine) Professor of  
Radiology, Harvard Medical School; Director of Joint Program in  
Nuclear Medicine at Beth Israel Hospital, Brigham and Women's  
Hospital, Children's Hospital and Institute, and Dana Farber Cancer  
Center; Vice-President, National Commission on Radiological  
Protection and Measurements; 25 Shattuck St., Boston, MA 02115  
(617-732-1535)

H. J. DUNSTER, B.Sc., C.B. (health physics) Formerly Director  
National Radiological Protection Board (United Kingdom), Member,  
International Commission on Radiological Protection; Residence: 52  
Thames St., St. Ebbes, Oxford, OX1 1SU, United Kingdom  
(011-44- 865-251-716)

A. S. KUBO, Ph.D., MBA, P.E. (civil and nuclear engineering)  
Vice President, Technical Applications, The BDM Corp. 7915 Jones  
Branch Drive, McLean VA 22102 (703-848-7294)

H. G. PARETZKE, M.Sc., Ph.D. (radiation risk analysis) Head, Radiation  
Risk Analysis Section, GSF Institut für Strahlenschutz (Institute  
for Radiation Protection), Ingolstädter Landstrasse 1, D-8042,  
Neuherberg 2225 Federal Republic of Germany GE-055  
(011-49-893-187-2225)

F. L. PETERSON, Ph.D. (hydrology and geology) Professor of  
Hydrology and Chairman, Dept. of Geology and Geophysics, University  
of Hawaii, Honolulu, HI 96822 (808-948-7897)

W. J. SCHULL, Ph.D. (epidemiology: cancer, genetics, birth defects)  
Director of Center for Demographic and Population Genetics and  
Professor of Human Genetics, Univ. of Texas Health Science Center at  
Houston; Formerly Director of the Radiation Research Foundation at  
Hiroshima-Nagasaki, Japan. Address: Population Genetics, P.O.  
Box 20334, Houston TX 77225 (713-792-4680)

E. L. STONE, Ph.D. (soil science) Pack Professor Emeritus of Forest Soils, Cornell University; Adjunct Professor, Dept. of Soil Science, 2169 McCarty Hall, Univ. of Florida, Gainesville, FL 32611 (904-392-1956)

Consultants nominated by the Rongelap people

ROSALIE BERTELL, Ph.D., G.N.S.H. (biometrician) Editor in Chief, International Perspectives in Public Health; Commissioner International Commission of Health Professionals, Geneva; President, International Institute of Concern for Public Health, 830 Bathurst St., Toronto, Ontario M5R-3G1 Canada (416-533-7351)

UTE BOIKAT, M.Sc., Ph.D. (radioecology), Executive of the Department of Public Health, Freie und Hansestadt Hamburg, Tesdorpfstr.8, D-2000 Hamburg 13, Federal Republic of Germany. ((011-49)40-44195334).

BERND FRANKE, M.Sc. (radioecology), Executive Director (Washington Office), Institute for Energy and Environmental Research, 6935 Laurel Ave., Takoma Park, MD 20912 (301-270-5500)

Others who have informally helped in the production of this report:

## 2. BACKGROUND -- THE RONGELAP EXPERIENCE

Rongelap Atoll is located about 2,500 miles southwest of Hawaii, at 12°N, 167°E (Fig. 2 #1). It comprises more than 50 low-lying islands and islets, total area 3.07 sq. miles, which bound a lagoon of 400 sq. miles. The largest and by far the most important island, Rongelap, has an area of 0.3 sq. miles.

The geological structure is that of a coral reef atoll resting on a submerged volcanic mass. The islands are made of reef debris, primarily of sand and gravel size, and reef organisms.

The atoll is typical in appearance, and the islands are covered with vegetation. However, a major factor limiting the kinds of plants that can be grown as staples is the long dry season.

The Marshall Islands Statistical Abstract of 1986, issued by the Republic, lists the population of the atoll as totalling 235. Previously, it was 165 in 1973, 189 in 1967, 264 in 1958. In 1954 at the time of the Bravo incident, 84 persons were evacuated. (These fluctuations reflect the need to work elsewhere.) Earlier records for Japanese and German periods of control are: 99 in 1945, 98 in 1935, 110 in 1920, 100 in 1906, 120 in 1860.

However, Mr. Peter Oliver, the Republic's Special Assistant for Compact Affairs, has informed me that the Rongelap Distribution Authority now makes per capita payments from its Nuclear Claims Fund to 1,578 individuals. Currently, these amount to \$1480 per year to those exposed to fallout in 1954, and \$480 to others. The Council has also determined that 2,277 individuals qualify for the benefits of the Section 177 Health Care Program as a result of their ties to Rongelap.

### 2.1 Bravo test -- 1954

The initial event occurred on March 1, 1954, when a 17-megaton-yield thermonuclear device was set off at Bikini Atoll, the Bravo test. The device was 1000 times as powerful as the bombs that destroyed Nagasaki and Hiroshima; its cloud rose 25 miles above the earth, and after 10 minutes had a diameter of 70 miles.

It had been planned that the "cloud" would be blown to the west and north (Fig. 2.1 #1). Unexpectedly for whatever reason (Note 3), it was blown to the east so that at about 5 hours after detonation fallout began at Rongelap Atoll, and during the ensuing 7 hours fell in such quantities as to suggest to Rongelapese, who had never seen snow, that it was snowing (Sharp & Chapman, 1957). Rather than avoiding contact, children played in the powdery, finely granular fallout, and no particular effort was made to separate it from food or clothing. No warning was or had been issued by the military.





165°E

180°

165°E

150°E

PACIFIC  
OCEAN



Copyright 1982 by  
Continental Air Micronesia  
Route Map, Actual flight path

Fig. 2.1 #1 LOCATION OF THE MARSHALL ISLANDS

About 50 hours after the "shot", the Navy removed the 64 Rongelap residents from the Atoll to the medical base at Kwajalein (Sharr & Chapman, 1957; Cronkite et al, 1956). Also, eighteen visiting Rongelapese were removed from Sifo Island, Ailingnae Atoll, and 157 Utirik people from Utirik Atoll. It was immediately recognized that the surveillance and care of these people required far more professional staff than the base could supply, and a special medical team hurriedly organized for this purpose in the United States, utilizing naval and AEC personnel, reached the base 8 days after the detonation.

Consistent with a whole-body dose of 190 rem (over two days), two-thirds of the Rongelap group experienced nausea, 10% with vomiting and diarrhea, which cleared within three days or so, and all showed depressed white-blood-cell counts (Cronkite et al, 1956). As a result of the skin dose from physical contact with fallout, about 70% developed skin lesions of widely varying severity after a latency period of two to three weeks. Most of these were to heal successfully but a few developed significant scarring.

The most "significant" part of the initial exposure produced no immediate signs or symptoms. A half-dozen thyroid-seeking radionuclides entered the body through fallout-contamination of food and water. Over the course of the following weeks these iodine and tellurium radionuclides delivered doses that eventually caused thyroid hypofunction and the appearance of thyroid tumors.

The Bravo test posed new dosimetry problems, only vaguely sensed before. Owing to the gigantic energy-yield at ground level, great quantities of coralloid radioactive material were generated (Hiroshima and Nagasaki had involved high air-bursts): 142 radionuclides were involved whose radiations and rates of decay varied greatly, and whose eventual effects depended on the weather conditions and the living habits of the exposed population.

At the time of evacuation, the exposure rate in Rongelap village was 1.2 - 2.3 R/hour. The whole-body dose of "175 R in air" reported in 1956 was approximately correct. The dose estimate for the thyroid gland, however, was much too low because only iodine-131 had been considered in the calculation. As a result, the appearance of thyroid disease later on was quite unexpected.

An upwards revision of thyroid dose was reported in 1964 when iodine-133 and iodine-135 were included. (James, 1964). The revisions of 1984 (Lessard et al, 1985; Lessard, 1984a), based on a comprehensively planned attack on the problem (Bond et al, 1978), put the mean adult whole-body dose at 190 rem. The revised total dose to the thyroid gland, including contributions from all seven important radionuclides was greatly increased and varied significantly with age at exposure in 1954 -- from 5,200 rem for a one-year old to 1,600 rem at age 14, and 1,200 rem for the adult male. It was estimated that 95% of the thyroid dose was received during the first three post-exposure weeks, and 100% within three months (Note 4).

1964-75. Unquestionable damage to the thyroid gland, especially to those exposed below the age of 10, made its appearance. A reexamination of earlier estimates of dose to the thyroid gland led to their elevation by a factor of about 2 for adults, and 5 or more for children. The administration of thyroid hormone (interrupted on occasion) to the entire exposed population was begun in 1965 as a prophylactic measure against thyroid neoplasia (nodules, cancer), and also to correct for possible losses in thyroid function.

By the end of 1974 (Fig 2.3 # 1), the thyroid tumor record was as follows:

Age below 10 in 1954: 17 tumors in 19 persons examined, including 1 cancer.

Age 10-18 years in 1954: 2 tumors in 12 persons examined.

Age above 18 years in 1954 : 3 tumors in 33 persons examined, including 2 cancers.

Almost all persons with thyroid nodules were sent for surgical treatment to the Cleveland Metropolitan Hospital, Cleveland, Ohio. Each one was compensated at the rate of \$25,000 per surgery.

The occurrence of thyroid disease as well as a case of acute leukemia worried the Rongelap people. The medical team was accused of having deceived the Rongelap people and of using them as guinea pigs. The Brookhaven medical services were boycotted during 1972, but they were accepted later in the year after a favorable report on the matter by an international committee.

1976-79. More thyroid nodules appeared. The Rongelap people continued to be worried. They asked for an independent health review which was not granted. A group of Brookhaven scientists proposed a comprehensive dosimetry review (Bond et al, 1978), which DOE then funded (Lessard, 1984a; Lessard et al, 1984c; Lessard et al, 1985). Independently, DOE initiated a "Northern Marshall's Survey" based on an aerial survey by EG&G and some terrestrial work by Lawrence Livermore National Laboratory (Robison et al, 1980; Robison et al, 1982b; Tipton & Meibaum, 1981).

1980-84. DOE summarized its survey results in 1982 with a report in Marshallese, embellished with colored illustrations. (This is the DOE-1982 book under review in the present report. See Note 1.) The conclusion, that Rongelap Island was safe, was not accepted by all of the people. The Rongelap people requested the Government to transfer them to another atoll. Significant parts of the anti-nuclear documentary film, Half-Life, were filmed at Rongelap. The film suggested that the people had been used as "guinea pigs".

1985. The Rongelap people abandoned Rongelap and sailed for Majiето Island in Kwajalein Atoll. The U. S. Congress passed the Compact of Free Association Act of 1985 (Public Law 99-239) of which Section 103(i) is the basis for the present inquiry (Note 2).

1987 The following points are of major interest for the present report.

(a) A clear distinction should be made between the late effects of the large acute exposure in 1954 (190 rem whole-body) and the possible (but as yet undetermined) effects of the much smaller chronic dose since resettlement in 1957 ( 3.5 rem or less to 1978).

(b) The original dose estimates for the 1954 exposure were much too low for the thyroid gland (Cronkite, 1954; Dunning, 1957). The necessity for major correction later on weakened or destroyed Rongelap confidence in DOE. The residual radiation doses during the first years of resettlement may also have been underestimated, but the corrections would be very much smaller.

(c) The occurrence of thyroid tumors ( ~ 30%) 10 years or later after returning to Rongelap (Fig. 2.3 #1; Note 4B) has been a confusing experience for the Rongelap people. In addition, eight cases of hypothyroidism have been observed (Adams 1988).

(d) No significant increase in tumors outside of the thyroid gland has been seen (Adams et al, 1984), except for 1 basal cell epithelioma in 1987 (Adams 1988) in the 81 persons at risk.

(e) No obvious gross difference in survivorship between 1954-exposed and 1954-unexposed groups has occurred (Fig. 2.3 #2). Although statistically significant decreases in some blood-cell types have been noted (Adams et al, 1982), none has been clinically significant.

(f) Based on four parameters (longevity, thyroid nodules, carcinoma, blood counts), there is no evidence of effects from the chronic low-level exposure associated with length of residence on Rongelap since 1957 (Note 4(b)). These studies are admittedly exploratory and cover only a small part of the health spectrum. However, the average dose over the period 1957-78 is quite small (3.5 rem or less), and will be accumulated at lower rates in the future.






FIGURE 2.3 #2 Survival as a function of time after 1954.

The numbers exposed and whole-body doses were: Rongelap, 67 persons, 190 rem; Ailingnae, 19 persons, 110 rem; Utirik, 167 persons, 11 rem. The unexposed group of 86 Rongelapese was matched (age, sex) in 1957 to the Rongelap-Ailingnae group and has been followed for survival annually.

(Figure courtesy of W. H. Adams, Brookhaven National Laboratory.)

### 3. REASSESSMENT

With the foregoing as background, let us now attempt to answer the questions which the Congress has asked: Were the doses calculated by DOE for 1978 correct? Does it follow that Rongelap is habitable? If not, what should be done?

It should be noted that the technical position has changed since 1982. More data have been published so that the original meager sampling has become more robust. In addition, we shall consider the findings of the Brookhaven National Laboratory, using an important method which DOE-1982 failed to consider, and also our own findings.\*

The data base employed by DOE-1982 comprised the results of the Northern Marshall Islands Survey of 1978 (September-November) which had been planned as an aerial reconnaissance to map external gamma-ray exposure rates (normalized to 1 meter above ground level) (Tipton & Meibaum, 1981). Two helicopters were employed, operating from a major support vessel, the U.S.N.S. Wheeling.

Subsequently the Livermore Laboratory program was added to obtain soil, water, vegetation and fish samples at each atoll "as time and facilities might permit" (Robison et al, 1982, Part 1). The time spent at Rongelap Atoll permitted 7 days for 9 islands, of which the major one was Rongelap. Operating from a large ship that had to cruise at a considerable distance offshore, and whose primary function was aerial reconnaissance, restricted the terrestrial work significantly.

The radionuclides dealt with were five: cesium-137, which is distributed throughout the body; strontium-90, a bone seeker; plutonium-239.-240 and americium-241, which have very long half-lives and which are tightly bound by bone, liver and testes (Table 3 #1).

The Livermore group took soil samples from some 20 scattered locations on Rongelap Island whose averages (picocuries/gram) for 0-10 cm depth were: cesium-137, 12; strontium-90, 7.1; plutonium-239,-240, 2.6; americium-241, 0.9 (Table 3 #2).

This soil contamination provided the basis for human exposure in two ways. Radiations emanated from the ground or standing vegetation leading to external dose. Radiations that emanated from food and water after entering the human body were responsible for internal dose.

---

\* B. Franke states that the enabling legislation calls for study of only the original findings and report. A second committee should consider subsequent findings, and a third group should execute its recommendations.

The total dose received was the sum of the external and internal doses. The external whole-body dose was estimated by measuring the exposure in air (e.g., at 1 meter above ground) and applying a factor based ultimately on measurements with phantoms to the meter reading. The internal dose was estimated by the Livermore group on the basis of an assumed diet and the analysis of the radionuclide contents of Rongelap food products in it.

The lagoon and its fish were found to be a trivial source of dose. Ground water (well water) was an unimportant source, since its activity was very low and, in any case, the people relied heavily on catchment of rain rather than wells (Noshkin et al 1981).

Before considering the data, the nonprofessional reader may wish to consult Note 6 which explains the radiological usage of such terms as exposure and dose, and the definition of their units. It may also be noted here that my use of the term whole-body dose (internal) usually signifies the committed effective dose equivalent; the tissue dose (internal) is usually the committed dose equivalent. The Livermore Laboratory calculated its doses as integral doses, i.e., for a stated period of time, the annual dose for each year was summed.



TABLE 3 #1 SOURCES OF FALLOUT RADIATION AT RONGELAP

Radionuclide	Half-life <sup>a/</sup>	Principal radiations <sup>a/</sup>			ICRP-derived limit on daily oral intake <sup>e/</sup>	Fraction absorbed from gut <sup>f/</sup> in adults
		$\alpha$ <sup>b/</sup>	$\beta$ <sup>c/</sup>	$\gamma$ <sup>c,d/</sup>		
	years	MeV	MeV	MeV	pCi/d <sup>f/</sup>	
Cesium-137	30	-	0.187	.66	9860 * 5920 **	1.0
Strontium-90	29	-	1.13	-	2470 * 1480 **	.3
Plutonium-239	24,065	5.23	-	-	30 ** (60)	.001
-240	6,537	5.24	-	-	30 ** (60)	.001
Americium-241	432	5.57	-	-	37 ** (67)	.001

a/ ICRP Publication 38. (Radionuclide transformations)

b/ Quality factor, 20

c/ Quality factor, 1

d/ X and gamma rays are omitted whose total contribution to dose would be less than 10%.

e/ Derived from ICRP Publications 30 and 48. The ICRP limit on intake for workers was divided by 30 (\*) to bring the annual committed effective dose-equivalent to 170 mrem, or by 50 (\*\*) for 100 mrem. The ICRP limit includes a factor of 2 to prevent any one tissue receiving more than 50 rem. That factor is unnecessary in the present low-dosage case. The numbers in parentheses give the applicable guide without such correction.\*

f/ ICRP Publication 30. Supplement to Part 1. (Annals, Vol. 3), and ICRP Publication 48 for transuranics.

\*John Dunster adds: The intake limits apply to adults. For children, the strontium limit should be divided by a factor of about 3, and those for plutonium and americium by about 2. (National Radiation Protection Board G 87, Aug 87.)

TABLE 3 #2

RONGELAP ISLAND: RADIONUCLIDE SOIL PROFILES<sup>a/b/</sup>

Depth (cm)		Average specific activity for dry soil (pCi/g)							
		Cesium-137		Strontium -90		Plutonium -239,-240		Americium -241	
		1978	1987	1978	1987	1978	1987	1978	1987
0-5	0-10	15	10.6(7)	6.9		3.2		1.0	1.7(3)
5-10		9		7.7		2.0		.78	
10-15	10-20	5.4		6.7		1.1		.41	
15-25		2.6		4.5		.35		.18	
25-40		1.8		2.1		.07		.08	
0-40		5.0		4.6		.89		.35	
Number of profiles		27		20		18		.7	

<sup>a/</sup> The 1978 profiles are from Robison et al, 1982, Part 4, Appendix B.

<sup>b/</sup> The 1987 values are from Boikat and Paretzke (Note 8). The number of samples is given in parentheses. They are corrected back to 1978.

#### 4. DOSE

DOE-1982 reported three doses for the Rongelap people who would live on Rongelap Island for the period 1978-2008, tacitly assuming a constant diet. To this DOE-1982 added the stipulation that the diet would be based on "local food only from Rongelap Island" (Note 1).

It should be pointed out, however, that the stipulation of "local food only" is incorrect. The doses used by DOE-1982 were estimated by Robison et al (1982b), who based them on the type B community diet described by Naidu et al (1980). That diet involves imported foods brought in on a regular basis by supply ship.

The three doses are as follows:

(1) The "highest average amount of radiation the people might receive in any part of the body" was 2.5 rem. I take this to be Livermore's "integral dose" in which each year's delivery is summed over 30 years (Robison et al, 1982b, Table 17). I will compare it to the committed whole-body dose (rem) over 30 years (i.e., the committed effective dose equivalent for a standard man).

(2) The corresponding bone marrow average would be 3.3 rem (Robison et al, 1982b, Table 14). I take this to be the "tissue dose" and it is approximately equal to the committed dose equivalent.

(3) The highest dose to any one person was set at 0.4 rem this being three times the average dose.

For orientation, it may be said that DOE's whole-body and bone-marrow doses are for practical purposes confirmed by recalculations employing the original data and corrected assumptions, and by those employing subsequent findings on additional field samplings.

However, the independent assessment by the Brookhaven National Laboratory, based on whole-body counting for cesium and urinary analysis for strontium, lowers the whole-body dose significantly. This estimate, in my opinion, is the definitive one.

Brookhaven's estimate of the transuranic dose (plutonium, americium) has raised the question of the size of its contribution to dose--a matter which is under discussion--but in any case, apparently not great enough to prevent a decision from being made. This matter will be discussed.

The question of infant dosage, neglected previously, has been dealt with specifically (or will be).

#### 4.1 External Dose

The aerial survey (Tipton & Meibaum, 1981) provided DOE with important information on exposure to fallout in the Northern Marshall Islands. As the survey proceeded south and east from Bikini Atoll, the seat of the Bravo shot, the external exposure rate fell ( Table 4.1 #1). It was calculated for 1 meter above ground level.

At Rongelap Atoll (Figure 4.1 #1), the islands fell into four exposure groups (microreoeotgens per hour) from north to south: Naen, Yugui, Lomuila (28-43  $\mu$ R/h), Eniaetok, Kabelle, Gogan (10-27  $\mu$ R/h); Busch, Borukka, Gabelle, Tufa (5-9  $\mu$ R/h); Rongelap and Arbar (4.1-4.5  $\mu$ R/h).

The external dose (whole-body), was calculated from exposure by my assuming 1 roentgen = 0.7 rem (Kerr, 1980). For Rongelap Island the annual dose was .028 rem, well below the EPA guide of .170 rem/year; 8 other major islands were also below the guide (Table 4.1 #1).

There is also a shallow dose to be considered, that due to beta rays which travel for short distances into those parts of the body that are near or in close contact with the soil and that are unshielded. Their contribution is considered to be negligible (Note 9).

These estimated external gamma-ray dose rates are maximal ones. Indoors the rate is reduced by about 50%. Likewise, the rate is reduced by about 50% in the immediate vicinity of houses owing to the coral gravel that is spread around them (Shingleton et al, 1987 and Robison et al, 1982b).

Other annual contributions to external dosage which are not included come from cosmic radiation (.028 rem) and medical exposure.

In summary, the contribution of fallout to the total external radiation dose at Rongelap Island in 1978 was approximately .028 rem per year uncorrected for the shielding within or around buildings, which would decrease it by 25% or more. The 30-year whole-body dose would be .590 rem allowing for spontaneous decay, but not shielding. Environmental decay such as leaching of radionuclides from the soil would reduce this estimate still more, but was not allowed for.


RONGELAP ATOLL

Figure 4.#1 PRINCIPAL ISLANDS OF RONGELAP ATOLL

The numbers in parentheses are the external whole-body exposure-rates in microroentgens/hour, corrected for cosmic radiation, as determined in 1978 by aerial survey (Tipton & Meibaum, 1981).

TABLE 4.1 #1 AVERAGE EXTERNAL EXPOSURE AND EXTERNAL DOSE RATES  
(gamma ray) FOR ISLANDS AFFECTED BY BRAVO FALLOUT

Atoll and Reference	Island	Year	a/ Exposure (gamma)	b/ Dose (whole-body)
			microroent- gens/hour	rem/year
<u>Bikini Atoll</u>				
Tipton & Meibaum (1981)	Eneu	1978	2.7	.017
	Bikini		35.0	.215
Shingleton et al (1987)	Eneu	1986	--	.018
	Bikini		--	.160
<u>Rongelap Atoll</u>				
Tipton & Meibaum (1981)	Rongelap	1978	4.5	.028
	Arbar		4.1	.025
	Busch, Tufa, Borukka, Gabelle		5-9	.031-.055
	Eniaetok, Kabelle, Gogan		10-27	.061-.166
	Lukuen, Naen, Yugui, Lomuila		28-43	.172-.264
Paretzke (Note 8)	Rongelap	1987	4.1 (7) <sup>cd</sup>	.025
Greenhouse & Milten- berger (1977)	Rongelap	1977	3.6-4.5	.022-.028
<u>Ailingnae Atoll</u>				
Tipton & Meibaum (1981)	Sifo	1978	1.4	.009
Paretzke (Note 8)	Mogiri	1987 <sup>d</sup>	1.3 (1)	.008
	Enibuk		2.2 (1)	.013
<u>Utirik Atoll</u>				
Tipton & Meibaum (1981)	Utirik	1978	0.8	.005

a/  
Measured at 1 meter above ground level, corrected for cosmic rays.

b/  
Annual, whole-body dose (millirem/year) calculated as equal to  
6.13 x  $\mu$ R/hour. For the epidermal dose, see Note 9.

c/  
The average of 7 locations ranging from 2.2 to 4.6  $\mu$ R/hour.

d/  
Corrected for decay back to 1978.

#### 4.2 Internal Dose - Lawrence Livermore National Laboratory

Lawrence Livermore attacked the problem by determining what went into the body by ingestion and inhalation (picocuries per day), and then applying appropriate factors to such input (exposure) to obtain the dose in rem. The particular ones I have used are given in Table 4.2 #1.

Ingestion. The major uncertainty lies in the diet--no one knows precisely what it is, although several attempts have been made to define it. To be on the safe side, DOE-1982 chose the BNL community B diet, i.e., one involving a greater amount of food and also a greater input of contaminated food (Note 11). Naidu et al (1980) who originally described it commented that the diet represented prepared, not eaten food, and that in fact it was more than a person could eat. This results in overestimation of dose. The Lawrence Livermore group that used it for dose calculations concurred.

The 1978 specific activities measured by the Livermore team were made on 21 samples of coconut, 5 of Pandanus, 1 of breadfruit, 1 chicken, 2 pigs and 98 fish, on the whole a barely adequate number (Robison et al, 1981a, 1982b). In 1986, however, that Laboratory took additional samples (Robison 1988), and in 1987 this reassessment project also collected some which were analyzed independently. The results, summarized in Table 4.2 #2, show remarkable agreement for the Livermore 1978 and 1986 cesium data on the foods contributing the major part of exposure and also good agreement for our independent samples in 1987 (Note 8).

I am therefore taking 4400 picocuries/day as the exposure due to cesium-137, based on a total of about 4000 for foods listed in Table 4.2#2 plus a 10% allowance for a miscellaneous variety of others (Note 11, Table #1). The whole-body, red marrow and bone surface doses for 30 years are just about equal, 1.65 rem (Table 4.2 #1).

The strontium estimates at present are based on the original 1978 sampling. (No strontium analyses were done on the Livermore 1986 samples, nor were our 1987 samples delivered soon enough to have them done on time.) I am therefore taking .035 picocuries/day for the exposure, based on the field samples plus a 25% increment for other miscellaneous foods. The 30-year doses for whole-body, red marrow, and bone surface are .032, .175 and .385 rem, respectively.

In the case of the transuranics, the Livermore group is now summarizing their Rongelap work through 1987 and this involves some revision of both data and dose calculations (Table 4.2#3). Based on a

TABLE 4.2 #1A

INGESTION

FACTORS TO CONVERT "INITIAL DAILY INTAKE (pCi/d)" TO

"WHOLE BODY" OR "TISSUE" DOSE (rem) FOR DIFFERENT PERIODS OF DAILY INTAKE <sup>a/</sup>

Radionuclide & period	C.E.D.E. <sup>b/</sup>	Red marrow	Lungs	Bone surfaces	Liver
<u>CESIUM-137</u>					
initial year	1.7 E-5 <sup>c/</sup>	1.8 E-5	Like C.E.D.E		
0-30 year	3.7 E-4	3.8 E-5			
30-70 year	2.2 E-4	2.4 E-5			
<u>STRONTIUM-90</u>					
initial year	4.7 E-5	2.4 E-4	1.8 E-6	5.3 E-4	1.8 E-6
0-30 year	9.2 E-4	5.0 E-3	3.6 E-5	1.1 E-2	3.6 E-5
30-70 year	5.6 E-4	3.0 E-3	2.2 E-5	6.6 E-3	2.2 E-5
<u>PLUTONIUM-239.-240</u>					
initial year	1.3 E-3	1.9 E-3	1.0 E-8	2.4 E-2	4.2 E-3
0-30 year	3.9 E-2	5.7 E-2	3.1 E-7	7.3 E-1	1.3 E-1
30-70 year	5.1 E-2	7.4 E-2	4.1 E-7	9.6 E-1	1.7 E-1
<u>AMERICIUM-241</u>					
initial year	1.3 E-3	Like plutonium			
0-30 year	3.9 E-2	5.7 E-2	1.6 E-6	7.3 E-1	1.3 E-1
30-70 year		Like plutonium			

<sup>a/</sup> It is assumed that the daily diet remains constant, but that the radionuclides in it decay spontaneously. The table provides dose factors in rem/picocuries/day. It is based on NRPB (1987) which provides factors in Sv/Bq (= 3.7 x rem/picocurie), and is consistent with ICRP recommendations (ICRP 1986, 1987). These factors allow for the fraction of radionuclide absorbed from the gut, its distribution and residence time in the body, the absorption and effectiveness of its radiation in the body, and its rate of physical decay.

<sup>b/</sup> Committed effective dose equivalent (whole-body dose). Other doses are committed dose equivalents (tissue dose). The C.E.D.E. is the sum of the dose equivalents to all tissues of the body of a standard man, each weighted by the risk resulting from a unit dose to that tissue as compared to the risk from a unit dose to the whole body.

<sup>c/</sup> E-5 signifies:  $\times 10^{-5}$ .



TABLE 4.2 #1B

INHALATION

FACTORS TO CONVERT "INITIAL DAILY INTAKE (pCi/d)" TO

"WHOLE BODY" OR "TISSUE" DOSE (rem) FOR DIFFERENT PERIODS OF DAILY INTAKE <sup>a/</sup>

Radionuclide & period	C.E.D.E. <sup>b/</sup>	Red marrow	Lungs	Bone surfaces	Liver
<u>CESIUM-137</u>					
initial year	1.0 E-5 <sup>c/</sup>	9.9 E-6	1.1 E-5	9.4 E-6	1.0 E-5
0-30 year	2.2 E-4	2.0 E-5	2.2 E-4	2.0 E-4	2.2 E-4
30-70 year					
<u>STRONTIUM-90</u>					
initial year	7.7 E-5	4.2 E-4	4.6 E-6	9.2 E-4	3.1 E-6
0-30 year	1.6 E-3	8.7 E-3	9.5 E-5	1.9 E-2	6.4 E-5
30-70 year					
<u>PLUTONIUM-239.-240</u> <u>&amp; AMERICIUM-241</u>					
initial year	1.5 E-1	2.3 E-1	2.3 E-2	2.8 E-0	5. E-1
0-30 year	4.5 E-0	6.9 E-0	6.9 E-1	8.4 E-1	1.5 E-1
30-70 year	6.0 E-0	9.2 E-0	9.2 E-1	1.12 E-2	2.0 E-1

<sup>a/</sup> It is assumed that the daily diet remains constant, but that the radionuclides in it decay spontaneously. The table provides dose factors in rem/picocuries/day. It is based on NRPB (1987) which provides factors in Sv/Bq (= 3.7 x rem/picocurie), and is consistent with ICRP recommendations (ICRP 1986, 1987). These factors allow for the fraction of radionuclide absorbed from the gut, its distribution and residence time in the body, the absorption and effectiveness of its radiation in the body, and its rate of physical decay.

<sup>b/</sup> Committed effective dose equivalent (whole-body dose). Other doses are committed dose equivalents (tissue dose). The C.E.D.E. is the sum of the dose equivalents to 11 tissues of the body of a standard man, each weighted by the risk resulting from a unit dose to that tissue as compared to the risk from a unit dose to the whole body.

<sup>c/</sup> E-5 signifies:  $\times 10^{-5}$ .

Table 4.2 #2 COMPARISON OF ACTIVITY MEASUREMENTS (Referred to 1978)  
CESIUM-137 and STRONTIUM-90

		a/b/ Livermore (collected in 1978)			b/ Livermore (collected in 1986)			b/c/ This Report (collected in 1987)	
Item	grams/ day eaten	# samples	pCi/ gram (fresh)	pCi/ day	# samples	pCi/ gram (fresh)	pCi/ day	pCi/ gram (fresh)	
<u>CESIUM-137</u>									
Copra nut products	293	(18)	6	1758	(4)	6.2	1817		
Drinking nut:									
Meat	100	(3)	2.6	260	(86)	2.3	230	(6)	4.3
Juice	514	(3)	1.4	720	(85)	1.3	668	(7)	1.6
Pandanus juice	96	(2)	11.1	1066	(26)	10.9	1046		
Breadfruit	36	(1)	2.7	97	(13)	3.4	122		
Pork	1.4	(2)	8.5	12		—			
Chicken	3	(1)	2.5	8		—			
Fish	194	(98)	.025	5		—			
Arrow root	0		0	0		—			
Coconut crab	1			?					
Limes									
TOTALS				3926			3883		
<u>STRONTIUM-90</u>									
Copra nut:									
Meat	168	(8)	.022	4					To be done
Juice	125	(10)	.004	0.5					
Drinking nut:									
Meat	100								
Juice	514	(3)	.0014	0.7					
Pandanus juice	96	(3)	.181 <sup>d/</sup>	17.4					
Breadfruit	36	(1)	.095	3.4					
Pork	1.4	(2)	.005*	0.1					
Chicken	3	(1)	.009*	0.1					
Fish	194	(98)	.01*	1.9					
Arrow root	0								
Coconut crab	1								
TOTALS				28.2					

a/ The activities with an asterisk are from Robison et al (1982b), the original report. The other specific activities are a personal communication from Dr. Robison and involve a revision of the original data.

b/ Number of samples in parentheses. A sample comprised 5-6 coconuts, 3-5 breadfruit, and 1-2 Pandanus fruits.

c/ See Note 8 for details. Well water: cesium-137, .03 pCi/liter; strontium-90, .03 pCi/liter; plutonium-239, .0024 pCi/liter.

d/ The fibrous part of the fruit has a 10-fold greater strontium content, but is not eaten. Cesium is the same in both parts.

TABLE 4.2 #3

PLUTONIUM-239,-240 AND AMERICIUM-240 IN 1978 FOODS  
AT RONGELAP ISLAND BASED ON BNL TYPE B DIET<sup>a/</sup>

Item	Grams per day	Plutonium -239.-240 pCi/gram-fresh	Americium -240 pCi/gram-fresh	Picocuries per day
Drinking coconut juice	514	$2.7 \times 10^{-5}$ (2)	$2.5 \times 10^{-5}$ (3)	.027
Copra nut products	293	$6.5 \times 10^{-5}$ (5-9)	$6.8 \times 10^{-5}$ (7-9)	.039
Pandanus juice	96	$6.0 \times 10^{-5}$ (5)	$2.7 \times 10^{-5}$ (3)	.008
Fish (reef)	194	$24 \times 10^{-5}$ ( 98)	$4.3 \times 10^{-5}$ ( 98)	.060

<sup>a/</sup> Livermore has revised the transuranic data of Robison et al (1982b), and the present doses are about 50% higher. The entries in the table above are based only on chemical determinations (number of samples in parentheses). They are responsible for about 25% of the total dose which Livermore now attributes to plutonium-239,-240 (.37 pCi/day) and americium-241 (.13 pCi/day). The rest of the dose was estimated by a ratio method of extrapolation: it was assumed that the Rongelap ratio, specific activity of food to that of soil (chemically determined) would equal the Bikini ratio (based on chemical determinations for both soil and food).

type-B-diet input of 0.5 picocuries/day (.37 pCi/d plutonium-239,-240 + .13 pCi/d americium-241), I estimate the following 30-year doses: whole-body, .020 rem; red marrow, .029 rem; liver, .065 rem; bone-surface, .365 rem. The Livermore doses are about a factor of 3 smaller, in large part because they are integral doses, not committed ones.

Inhalation. It is the transuranics that are of consequence. The original estimates of dust intake were very much too high (Shinn et al 1980) and they have been reduced to make them more realistic (Robison 1988). The daily intake for adults is estimated now at .0037 picocuries for plutonium-239,-240, and .0012 for americium-241. Their contribution to the effective whole-body dose would be about .023 rem in 30 years, and about 0.35 rem to the bone marrow, .075 rem to liver, and .42 rem to bone surface. The matter is discussed in Note 10.

Summary. Using the input method, the calculations of committed dose are in practical agreement with those of DOE-1982. It should be noted that these are for adults. It should also be noted that the estimates depend directly on the assumed diets. The following tabulation is a summary:

<u>30-year Dose (type B diet)</u>		
<u>Source</u>	<u>Whole-body dose</u>	<u>Red marrow dose</u>
	(rem)	(rem)
Inhalation	.023	.035
Internal doses:		
-cesium-137	1.63	1.67
-strontium-90	.032	.175
-transuranics	.02	.029
External dose	<u>.590</u>	<u>.590</u>
Totals	2.295	2.499
DOE-1982	2.500	3.300

For comparison, this project sampled three sites at Ailiniinae Atoll, which is not inhabited except for visits to gather food (Note 8). Landings were made on Mogiri, Gereia-Knox, and Enibuk Islands. The cesium-137 averages for the three sites for drinking-coconut meat and juice, and for the first 10 cm of soil, were 14% to 25% of the corresponding Rongelap averages. Two coconut crabs averaged 1.15 pCi/gram. The plutonium-239,-240 content was less than .006 pCi/gram.

#### 4.3 Internal Dose - Brookhaven National Laboratory

Brookhaven chose the method of whole-body counting to follow cesium in the exposed population, supplemented by urinary analysis to determine strontium and plutonium-239 (Conard et al, 1980; Lessard et al 1984b, 1984c; Miltenberger et al 1980). The method is the definitive one for cesium, since it is a direct measure of what is wanted and it is independent of assumptions regarding the diet and other external factors. It is of primary importance for the present case, since cesium accounts for some 80% of the internal whole-body dose.

The Brookhaven results in Fig. 4.3 #1 show the decline in cesium-137 body burden from about 670,000 picocuries in 1958-65 (.11 rem/year) to about 175,000 picocuries in 1979 (.03 rem/year). Thus the Brookhaven cesium internal dose-rate of .030 rem/year (whole-body) in 1978 was only 33% of that by the dietary input method (.094 rem/year). The 30-year cesium whole-body dose was .624 rem. The tissue doses to bone surfaces, red marrow, liver, etc. would be equal to this figure.

DOE-1982 overstated the cesium dose by a factor of three, relative to whole-body counting. The most likely source of the discrepancy would be the diet--the use of the type B diet. Robison (1983) has reported evidence that this could be so. If the MLSC diet (imports available) were employed (Note 11, Table 1), the cesium body content calculated from the input data (.19 microcuries) would be in approximate agreement for 1978 with that measured by whole-body counting (.17 microcuries). (Do Lessard and Robison agree to this statement?)

We do not have an independent field check on the accuracy of the whole-body field measurements. The point may be made, however, that it was this team that discovered the precipitous rise in body-burden of the Bikini settlers in 1977-78 and who therefore called for their removal from Bikini Atoll (Conard et al, 1980; Miltenberger et al, 1980).

In the case of strontium, we shall take the 1980 findings at face value. The annual whole-body dose based on urine analysis was about .001 rem, from which I calculate a 30-year dose of .021 rem. The corresponding tissue doses are: red marrow .11 rem; bone surfaces, .25 rem.




Figure 4.3 #1. Adult cesium-137 body burden as a function of time since resettlement of Rongelap Island in 1957.

The maintenance of the body content depends on the radionuclide intake from the diet. The physical half-life is 30 years; the physiological half-life is 110 days in men, 80 days in women, and less in youths and children. (1 Bequerel = 27 picocuries; 1 nanocurie = 1,000 picocuries) The maintenance of the specific activity of 1 pCi/g in soft tissue for 1 year gives rise to a dose of .01 rem.

(Figure courtesy of E.T. Lessard, Brookhaven National Laboratory.)

In the case of the transuranics, the background of the problem is worth mention. The quantity of plutonium-239 in the urine is minute, being something like  $.1$  to  $1 \times 10^{-3}$  picocuries/liter. It has only been during the past several years that the Brookhaven group has felt able to do accurate determinations using the new fission track method. More than 250 Rongelap samples have been analyzed, but none of these has been reviewed with respect to the history of the donor, i.e., age, period of residence on island, occupation, etc., owing to the fact that support for the project terminates this year.

At my request, to provide some orientation to this problem, the Brookhaven Laboratory gave Dr. Lessard the time for a brief survey. From a random sample of 35 determinations, the median urinary output was found to be about  $.03 \times 10^{-3}$  picocuries/day, equivalent to a dietary consumption of  $.13$  pCi/day (Note 7). However, the exceptionally broad distribution of the individual determinations calls for a detailed review which might reveal technical error, but could equally well point to hitherto unresolved or unsuspected physiological factors that influence the results.

The 30-year doses associated with a median urinary output of  $.03 \times 10^{-3}$  pCi/day of plutonium-239 are: whole-body,  $.0051$  rem; red marrow,  $.0074$  rem; bone surfaces,  $.092$  rem; liver,  $.017$  rem. The addition to these of the doses for plutonium-240 and of americium-241, which were not measured, would increase them by perhaps a factor of two.

The Brookhaven results may be summarized as follows:

	30-year dose 1978-2008*	
	Whole-body	Red marrow
Cesium-137:	.620 rem	.620 rem
Strontium-90:	.021	.110
Plutonium-239	.005	.007
Plutonium-240 } **	< .005	< .007
Americium-241 }		
External dose:	.59	.59
Total:	1.24 rem	1.33 rem

\* Not including inhalation

\*\* Estimated

The Brookhaven group summarized its results by calculating a 50-year dose from 1957 to 2008 (Lessard et al 1984c), based on a curve fitted to the observations from 1959 (?) to 1980, then extrapolating back to 1957 and forward to 2008 (Note 7, Tables #2 and #3). Adding up the annual doses thus obtained gives a total of  $.66$  rem (external + internal, but not including transuranics or inhalation).

#### 4.4 Infant Dosage

The following factors should be taken into account. The infant during the first six months may absorb from the gut a much greater fraction of radionuclide than the adult. The residence time of radionuclide in the body may be shorter than in the adult. For longer residence times, the amount retained is diluted by growth. The infant eats less than the adult.

In the case of cesium-137, which is completely absorbed from the gut in both infant and adult and whose residence time is short, the difference between adult and infant dose factors will be small. For plutonium-239, whose absorption by the infant is much greater and whose residence time is long, an appreciable difference can occur. However, because the transuranic contribution to the adult dose is so small, even if it be increased very appreciably in the infant, it will not necessarily be quantitatively important.

Balancing these variables against one another leads to the following committed dose factors (rem per picocurie daily intake) for whole-body exposure:

Radionuclide	<u>Factor at specified age (rem/pCi/day)</u>			
	0-1 yr	5 yr	10 yr	0-10 yr
Cesium-137				
Strontium-90				
Transuranics				



#### 4.5 Dose Summary

DOE-1982 stated the whole-body dose (integral) to be 2.5 rem for the period 1978-2008, of which 1.63 rem stems from cesium-137. That dose, based on the type B community diet, is about 1 rem too high for the following reasons.

Whole-body counting is the superior method for the determination of the cesium-137 whole-body dose. Based on 1978 conditions at Rongelap Island, the cesium dose by that method for 1978-2008 would be .62 rem (committed effective dose equivalent).

For strontium-90, the urine-derived dose of .021 rem is 60% of that calculated from the diet (.035 rem). The difference is in the same direction as that for cesium, and is small enough in absolute terms so that it will not materially affect the outcome one way or the other.

For plutonium-239, the estimates based on urine (median value) and diet are close enough for practical purposes (.005 rem and .009 rem, respectively; total transuranic, .010 and .020 rem respectively). However, as noted above, the wide spread of the urine data do call for further investigation.

I therefore conclude that the doses in Table 4.5 # 1 fall well within the present EPA guide for the general population of the U.S.A. (5 rem for 30 years, committed effective dose equivalent, standard man; I also take 30 rem in any one tissue except lens). They also satisfy the ICRP and NCRP guides (3 rem).

Whether or not these estimated doses guarantee that no one in any one year will exceed the individual guide of 0.5 rem, I cannot say. By and large that should be so.

The increase in cancer mortality resulting from the dosages of Table 4.5 #1 can be calculated as follows. Suppose that 500 persons were to live continuously on Rongelap Island for the period 1978-2008. On the average each would accumulate a committed dose (whole-body) of 1.25 rem over that 30-year period. For simplicity, I will assume that each receives the dose all at once. Then, taking an overall cancer mortality factor of  $5 \times 10^{-4}$  per rem (Shimizu et al, 1987; Preston and Pierce, 1987), I find the increment to be:

$$500 \times 1.25 \times 5 \times 10^{-4} = .31 \text{ extra cases.}$$

The factor for first generation genetic defects is smaller than that for cancer mortality (National Academy of Sciences, 1972; NCRP, 1987a), being approximately  $1 \times 10^{-4}$ .

The foregoing comments apply to the future. But what about the past? The Rongelap residents exposed to the Bravo shot received an acute dose of 190 rem in 1954 and during 1957-1978 they received a chronic dose of 1-3 rem. My opinion is that the addition to these past doses of something like 1.25 rem during the next 30 years will not appreciably increase detectable health and genetic risks in a way that should preclude return to Rongelap Island.

TABLE 4.5 #1

PROJECTED ADULT COMMITTED DOSES (1978-2008)  
FOR RESIDENCE ON RONGELAP ISLAND

Radionuclide	Whole-body <sup>a/</sup>	Red marrow <sup>b/</sup>	Bone surfaces <sup>b/</sup>	Liver <sup>b/</sup>
	rem	rem	rem	rem
<u>Internal:</u>				
Cesium-137	.62	.62	.62	.62
Strontium-90	.021	.110	.250	< .001
Transuranics <sup>b/</sup>	.010	.015	.184	.014
<u>External:</u>	<u>.59</u>	<u>.59</u>	<u>.59</u>	<u>.59</u>
Totals	1.24	1.32	1.64	1.24

<sup>a/</sup> Committed effective dose equivalent (standard man) = whole-body dose. The current guide in the U. S. is 5 rem in 30 years. The type B diet is assumed.

<sup>b/</sup> I would employ a guide of not more than 30 rem to any one tissue over 30 years, but due allowance must be made for the doses received by other tissues (ICRP No. 30).

<sup>c/</sup> Plutonium-239, -240 and americium-241.

## 5. DISCUSSION AND RECOMMENDATIONS

The conclusions reached and the issues raised by the body of this report are quite straightforward. The dose received is due to radiations from (a) soil and vegetation externally, and (b) from the food eaten. The review has shown that DOE-1982 overestimated the 1978-2008 adult dose at Rongelap Island. The whole-body dose reported now (1.25 rem, 30-year) is one-half of theirs; for the red marrow it is 40% (1.34 rem). Both sets of values (DOE-1982 and ours) are well below the current U.S. whole-body guide of 5 rem. I conclude that a return to residence on Rongelap Island is permissible.

(The doses in this report "start" in 1978. The current 1988 dose, 10 years later, would be about 20% less.)

### 5.1 Assumptions

Within the simple statement on return are several tacit assumptions. Living conditions on return should be equivalent to those prior to leaving in 1985. In particular, the diet should be equivalent to the former one and thus should meet the following conditions.

(a) The food consumed was in part raised locally, but was also purchased when the supply ship visited at regular intervals. I assume that as much money would be available now as was available then.

(b) In addition, the families received foods distributed by the USDA Special Food Assistance Program, but which has only one more year to go. In the final year, the allotment will be one-quarter of what it has been. I understand that a request for a 3 or 5 year extension is being asked for. The extent to which this program, or an equivalent one, could continue into the future will require discussion.

(c) I have been told that it was only in 1982 that the people became aware of the restriction on food gathering in the more northern islands (e.g., Naen). That restriction should remain in force.

(d) Looking at the map in Fig. 4 #1, one can see how the external exposure rate (i.e., that from soil and vegetation) increases on both sides of the lagoon as one goes from the southernmost islands of Rongelap and Arbar toward the north. For the time being I would consider as forbidden territory all islands to the north of Borukka and Eniaetok. All to the south are suitable for food gathering and residence.

(e) There are no restrictions on fishing, anywhere. Terrestrial crabs are restricted like other foods.

(f) There are no restrictions that apply to Ailingnae Atoll.

(g) I would also add to these restrictions that no arrow root be consumed. Little was consumed during the 10-15 years prior to leaving in 1985 because, as I understand it, there was none on Rongelap Island. Since then the plant has returned. The plant is troublesome to prepare, and I would suppose that as long as supplies of flour and rice are available, it will not be used.

## 5.2 Infant Dosage

To be done. This section may or may not be necessary.

## 5.3 Plutonium

Plutonium poses a special problem that has two facets. First, the dosage of plutonium calculated from the type B community diet does not agree with many individual estimates based on urinary excretion. Second, the determination of plutonium in the urine has been exceptionally variable from subject to subject. To represent this wide distribution I have used the median value (middle value), not the mean (average) value, of the entire group.

The problem should be approached from the perspective provided by the data in Table 4.5 #1. The transuranics (plutonium-239, -240 and americium-241) contributed less than 1.5% to the total whole-body dose. Suppose that they had been underestimated by a factor of 100. Their contribution would then rise to 1.6 rem, which added to the 1.25 rem from other sources would give a total of 2.85 rem. This dose is still within the guide.

As noted in Section 4.3, the great variations among the individual plutonium determinations do merit investigation and I urge DOE's support. I suggest that they are not entirely methodological, but stem from physiological variations due to age or other factors. It would be especially important to study the people before they return to Rongelap to determine how rapidly the body content is excreted and the relation of the excretion rate to various physiological factors, as well as after their return for purposes of monitoring.

Once the variation in the urine determinations is understood, their agreement or lack of agreement with the calculated output from an assumed diet could be attacked, so that the estimated dosages would become much more reliable.

I understand that DOE is now considering the matter.

## 5.4 Monitoring and Health Programs

I recommend that the whole-body counting program to determine cesium-137 should be resumed as soon as practical. (It was discontinued in 1985.) It should be supplemented at the same time by studies on the strontium and plutonium content of the urine. These studies are essential for the control of the population's exposure to the radionuclides that contaminate the atoll.

Carried out properly, such studies are also of prime interest to scientists throughout the world who are interested in preserving the health of people who have been exposed to nuclear radiations. I know that the Rongelap people do not want themselves to be "guinea pigs" to

satisfy the curiosity of research workers. But that is not the case here. The work done would help the Rongelap people themselves, and its results at the same time would also help others.

I expect the Rongelap people to receive routine medical care. But I would also expect certain groups of them to continue to be part of surveys for the appearance of cancer, to undergo blood tests that their physicians may consider to be important, and to help in providing accurate records of vital statistics. All of this cannot be done unless their physicians are allowed to examine them at regular intervals whether or not they feel ill.


### 5.5 Rehabilitation of Soil

After the Rongelap people have settled on Rongelap Island, a reexamination should be made of the levels of contamination at the other principal islands of the atoll, for the reasons given in Note 12. At present, the best estimate of their relative degrees of contamination is obtained from a comparison of the external exposure rates determined by aerial reconnaissance (Table 4.1 #1). Based on the results of the resurvey of the atoll and a consideration of the field trials at Bikini, a long-term plan should be drawn up.

The methods now available to combat the radionuclide contamination of soil are essentially two -- remove the upper layer of soil in which the contaminants concentrate, or treat the soil with potassium salts which block its uptake by plants. A variant of the latter is to wash the soil with sea water. A long-term plan might employ all three.


These methods have been under investigation at Bikini Atoll for some years (BARC 1987). Fig. 5.4 #1 illustrates for 4 coconut trees on Eneu Island (Bikini Atoll) how the application of potassium chloride to the soil decreased the contamination of the coconuts. Fig. 5.4 #2 illustrates the results for Bikini Island where the contamination is about ten times as great. Such treatment could be administered to islands of an intermediate level contamination in order to make them habitable. Their complete effectiveness against the highest levels, such as at Naen, is still under investigation, but a report on the matter should become available by next year.

Figure 5.5 #1



The  $^{137}\text{Cs}$  concentration in drinking coconut meat from 4 trees on Eneu Island (Experiment #2). The shaded area represents the time during which a total of 1800 lbs. of K per acre was applied. The "+" symbol represents the  $^{40}\text{K}$  concentration in the coconut meat. (This graph was supplied by Dr. W. L. Robison of the Lawrence Livermore National Laboratory.)

Figure 5.5 #2



The  $^{137}\text{Cs}$  concentration in drinking coconut meat from trees treated with 1000 lbs. of K per acre in 4 equal applications in 3 month intervals. The shaded area represents the time in which the K was applied. (This graph was supplied by Dr. W. L. Robison of the Lawrence Livermore National Laboratory.)



## REFERENCES

- Adams, W.H., J.A. Harper, R.S. Rittmaster, P.M. Heotis, W.A. Scott. (1982). Medical status of Marshallese accidentally exposed to 1954 Bravo fallout radiation: January 1980 through December 1982. BNL 51761 (Biology & Medicine-TIC 4500) (Available from National Technical Information Service.)
- Adams, W.H. (1985) Letter Report to U. S. Department of Energy.
- Adams, W.H. (1987) Personal communication to H. I. Kohn.
- Adams, W.H., J.R. Engle, J.A. Harper, R.S. Rittmaster, P.M. Heotis, W.A. Scott (1984). Medical status of Marshallese accidentally exposed to 1954 Bravo fallout radiation: January 1983 through December 1984. BNL 51958, Medical Dept., Brookhaven National Laboratory, Upton, NY 11973
- Bikini Atoll Rehabilitation Committee, (1987). Report No. 5, Status March 31, 1987. 1203 Shattuck Ave., Berkeley CA 94709
- Bond, V.P. et al. (1978). Surveillance of facilities and sites, dose reassessment for populations on Rongelap and Utirik following exposure to fallout. DOE Contract # EY-76-C-02-0016, 189# K-121
- Christy, M., R.W. Leggett, E.E. Dunning, K.F. Eckerman (1984). Age dependent dose conversion factors for selected bone-seeking radionuclides. ORNL/TM, 8929. Oak Ridge National Laboratory, Oak Ridge TN 37830
- Conard, R.A., L.M. Meyer, J.E. Rall, A. Lowery, S.A. Bach, B. Cannon, E.I. Carter, M. Eicher, H. Hechter (1958). March 1957 medical survey of Rongelap and Utirik people three years after exposure to radioactive fallout. BNL 501. Brookhaven National Laboratory, Upton, NY 11973
- Conard, R.A., L.M. Meyer, W.W. Sutow, A. Lowrey, B. Cannon, W.C. Moloney, A.C. Watne, R. E. Carter, A. Hicking, R. Hammerstrom, B. Bender, I. Lanwi, E. Riklon, J. Anjain. Medical survey of the people of Rongelap and Utirik Islands nine and ten years after exposure to fallout radiation (Mar. 1963 and Mar. 1964). BNL 908 (T-371) Medical Division, Brookhaven National Laboratory, Upton NY 11973
- Conard, R.A., K.D. Knudsen, B.M. Cobysn, et al (1975). A twenty-year review of medical findings in a Marshallese population accidentally exposed to radioactive fallout. BNL 50424, Brookhaven National Laboratory, Upton NY 11973
- Conard, R.A. et al (1980). Review of medical findings in a Marshallese population twenty-six years after accidental exposure to radioactive fallout. BNL 51261 (Biology & Medical TID-4500) Medical Dept., Brookhaven National Laboratory, Upton, NY 11973

11973

- ICRP (1979). International Commission on Radiological Protection. Limits for intakes of radionuclides by workers. ICRP Publication 30, Part 1. Pergamon Press, NY
- ICRP (1984). Principles for limiting exposure of the public to natural sources of radiation. ICRP Publication 39. Pergamon Press, NY
- ICRP (1985). Quantitative bases for developing a unified index of harm. ICRP Publication 45. Pergamon Press, NY
- ICRP (1986). The metabolism of plutonium and related elements. ICRP Publication 48. Pergamon Press, NY
- ICRP (1987). Data for use in protection against external radiation. ICRP Publication 51. Pergamon Press, NY
- James, R.A. (1964). Estimate of radiation dose to the thyroids of the Rongelap children following the Bravo event. UCRL 12273. Lawrence Livermore National Laboratory, Livermore CA 94550

- Kato, H., W.J. Schull, A. Awa, M. Akiyama, M. Otake. (1987). Dose response analyses among atomic bomb survivors exposed to low-level radiation. Health Physics 52: 645-52
- Kerr, G.D. (1980) A review of organ doses from isotropic fields of gamma rays. Health Physics 39, pp. 3-20.
- Lessard, E.T. (1988) Personal communication to Henry I. Kohn.
- Lessard, E.T. (1984a) Letter Report to Roger Ray, DOE Operations Office, P.O. Box 14100, Las Vegas, NV 89114
- Lessard, E.T., R.P. Miltenberger, S.H. Cohn, S.V. Musolino, R.A. Conard (1984c). Protracted exposure to fallout: the Rongelap and Utirik experience. Health Physics 46, 511-547
- Lessard, E.T., A.B. Brill, W.H. Adams (19 ). Thyroid cancer in the Marshall Islands: Relative risk of radioiodine and external radiation exposure. BNL 37232. Medical Dept., Brookhaven National Laboratory, Upton NY 11973
- Lessard, E.T., R. Miltenberger, R. Conard, S. Musolino, J. Naidu, A. Moorthy, C. Schopfer (1985). Thyroid absorbed dose for people at Rongelap, Utirik, and Sifo on March 1, 1954. BNL 51882. Brookhaven National Laboratory, Upton, NY 11973
- Lessard, E.T., X. Yihua, K.W. Skrable, G.E. Chabot, C.S. French, T.R. Labone, J.R. Johnson, D.R. Fisher, R. Belanger, J.L. Lipsztein. (1987). Interpretation of bioassay measurements. NUREG/OR-4884; BNL-NUREG-52063. Safety and Environmental Protection Division, Brookhaven National Laboratory, Upton, NY 11973
- Miltenberger, R.P., N.A. Greenhouse, E.T. Lessard (1980). Whole body counting results from 1974 to 1979 for Bikini Island residents. Health Physics 39: 395-407.
- Miltenberger, R.P., E.T. Lessard, J. Steiners, N.A. Greenhouse (1980?) <sup>137</sup>Cs in human milk and dose equivalent assessment. Undated manuscript given me by Lessard, Sept. 1987.
- Moss, W.D. (1988). Twenty-sixth Hanford Life Sciences Symposium, (October 1987): Modelling for scaling to man. (J.A. Mahaffey and J.A. McWhinney, co-chairmen). To be published as a special issue of the Journal of Health Physics.
- Naidu, J.R., N.A. Greenhouse, G. Knight, E.C. Craighead. (1980). Marshall Islands: A study of diet and living patterns. BNL 51313. Safety and Environmental Protection Division, Brookhaven National Laboratory, Upton. NY 11973
- National Academy of Sciences (1972). The effects on populations of exposure to low levels of ionizing radiation. Report of the Advisory Committee on the Biological Effects of Ionizing Radiations, Division of Medical Sciences, National Academy of Sciences, Washington D.C. 20006

- National Academy of Sciences (1980). The effects on populations of exposure to low levels of ionizing radiation. Report of the Advisory Committee on the Biological Effects of Ionizing Radiations, Division of Medical Sciences, National Academy of Sciences, Washington D.C. 20006
- NCRP (1957). National Council on Radiation Protection and Measurements Permissible dose from external sources of ionizing radiations. Insert to National Bureau of Standards Handbook 59, National Council of Radiation Protection and Measurements. 7910 Woodmont Av., Bethesda, MD 20814
- NCRP (1987a) Genetic effects from internally deposited radionuclides. NCRP Report No. 89. NCRP. 7910 Woodmont Av., Bethesda MD 20814
- NCRP (1987b) Recommendations on limits for exposure to ionizing radiation, NCRP Report No. 91. June 1, 1987. National Council on Radiation Protection and Measurements. 7910 Woodmont Av., Bethesda, MD 20814
- NCRP (1987c) Ionizing radiation exposure of the population of the United States. NCRP Report No. 93, National Council on Radiation Protection and Measurements, 7910 Woodmont Av., Bethesda MD 20814
- National Radiological Protection Board (1987). Interim guidance on the implications of recent revisions of risk estimates and the ICRP 1987 Coma statement. NCRP-GS9. Chilton, Didcot, Oxon OX11 0RQ, United Kingdom
- Noshkin, V.E., R.J. Eagle, K.M. Wong, T.A. Jokela, W.L. Robison (1981). Radionuclide concentrations and dose assessment of cistern water and groundwater at the Marshall Islands. UCRL-52853, Part 2. Lawrence Livermore National Laboratories, Livermore, CA 94550
- Preston, D.L., D.A. Pierce (1987). The effect of changes in dosimetry on cancer mortality risk estimates in the atomic bomb survivors. Radiation Effects Research Foundation Technical Report RERF TR 9-87.
- Robison, W.L. (1983) National Academy of Sciences conference.
- Robison, W.L. (1988) Personal communication to H. I. Kohn. These data should be published by LLNL in 1988.
- Robison, W.L., V.E. Noshkin, W.A. Phillips, R.J. Eagle (1980). The Northern Marshall Islands radiological survey: radionuclide concentrations in fish and clams and estimated doses via the marine pathway. UCRL-52853, Part 3. Lawrence Livermore National Laboratory, Livermore CA 94550
- Robison, W.L., C.L. Conrado, R.J. Eagle, M.L. Stuart (1981). The Northern Marshall Islands radiological survey: Sampling and analysis summary. UCRL 52853, Part 1. Lawrence Livermore National Laboratory, Livermore CA 94550

- Robison, W.L., M.E. Mount, W. A. Phillips, M.L. Stuart, S.E. Thompson, C.L. Conrado, A.C. Stoker (1982a). An updated radiological dose assessment of Bikini and Eneu Islands at Bikini Atoll. UCRL 53225, Lawrence Livermore National Laboratory, Livermore CA 94550
- Robison, W.L., M.E. Mount, W.A. Phillips, C.A. Conrado, M.L. Stuart, C.E. Stoiker (1982b). The northern Marshall Islands radiological survey: terrestrial food chain and total doses. UCRL 52858, Part 4. Lawrence Livermore National Laboratory, Livermore CA 94550
- Sharp, R., W.H. Chapman (1957). Exposure of Marshall Islanders and American military personnel to fallout. Operation Castle-Project 4.1 addendum. Armed Forces Special Weapons Project. Sandia Base, Albuquerque, NM. Document WT-938
- Shimizu, Y., H. Kato, W.J. Schull, S.L. Preston, S. Fujita, D. Pierce (1987). Life Span study report 11, Part 1. Comparison of risk coefficients for site-specific cancer mortality based on the DS86 and T65DR shielded kerma and organ doses. Radiation Effects Research Foundation RERF TR 12087
- Shingleton, K.L., J. L. Cate, M.G. Trent, W.L. Robison (1987). Bikini Atoll ionizing radiation survey, May 1985-May 1986. UCRL-53798. Lawrence Livermore National Laboratory, Livermore CA 94550
- Shinn, J.E., D.N. Homan, Robison, W.L. (1980). Resuspension studies at Bikini Atoll. UCID-18538, Lawrence Livermore National Laboratory, Livermore CA 94550
- Tipton, W.J., R.A. Meibaum (1981). An aerial radiological and photographic survey of eleven atolls and two islands within the northern Marshall Islands. EG&G Energy Measurements Group, Document EFF-1183-1758 (Available from the National Technical Information Service, Springfield VA 22161.)
- U. S. Congress. Compact of Free Association Act of 1985, U.S. Public Law 99-239, Section 103(i)
- U. S. Department of Energy. (1982). The meaning of radiation for those atolls in the Northern Part of the Marshall Islands that were surveyed in 1978. Washington, D.C.

## NOTES CITED IN THE TEXT

N.1

The following is quoted from "The Meaning of Radiation for Those Atolls in the Northern Part of the Marshall Islands That Were Surveyed in 1978", U. S. Department of Energy, Washington, D.C., November 1982, page 39:

**Information That Has Been Obtained from the Measurements  
Made in 1978**

If 233 people live on Rongelap Island and eat local food only from Rongelap Island

Scientists estimate that the largest amount of radiation a person might receive in one year from radioactive atoms that came from the U.S. bomb tests is 400 millirem. But usually the largest amount a person might receive would be less than this. This amount of radiation decreases every year, however, it decreases very slowly.

The highest average amount of radiation people might receive in the coming 30 years is 2500 millirem in any part of the body and 3300 millirem in just the bone marrow.

In the coming 30 years, scientists estimate that 10 people may die from cancers caused by things other than radiation from the atomic bomb tests. In addition to this, from 0.1 to 0.6 people may die in the future from cancers caused by radiation received in the coming 30 years from the atomic bomb tests.

In the coming 30 years, scientists estimate that 60 children could be born with health defects caused by things other than radiation from the atomic bomb tests. In addition to this, 0.007 to 0.1 children may eventually be born with health defects caused by radiation their parents receive in the coming 30 years from the atomic bomb tests.

If people live on Eniwetok and not on Rongelap Island, and eat local food only from Eniwetok, the amount of radiation they receive would be about the same.

If people go to Naen from Rongelap Island, and eat food from Naen, they might receive about five times more radiation while they are there.

If people go to Namu or Mele from Rongelap Island, and eat food from those two islands, they could receive about two times more radiation while they are there.

N-2

## COMPACT OF FREE ASSOCIATION ACT OF 1985

PUBLIC LAW 99-239—JAN. 14, 1986

99 STAT. 1783

department or agency of the United States or by contract with a United States firm) shall continue to provide special medical care and logistical support thereto for the remaining 174 members of the population of Rongelap and Utrik who were exposed to radiation resulting from the 1954 United States thermonuclear "Bravo" test, pursuant to Public Laws 95-134 and 96-205. Such medical care and its accompanying logistical support shall total \$22,500,000 over the first 11 years of the Compact.

91 Stat. 1159.

94 Stat. 84.

(2) **AGRICULTURAL AND FOOD PROGRAMS.**—Notwithstanding any other provision of law, upon the request of the Government of the Marshall Islands, for the first five years after the effective date of the Compact, the President (either through an appropriate department or agency of the United States or by contract with a United States firm) shall provide technical and other assistance—

President of U.S.

(A) without reimbursement, to continue the planting and agricultural maintenance program on Enewetak;

(B) without reimbursement, to continue the food programs of the Bikini and Enewetak people described in section 1(d) of Article II of the Subsidiary Agreement for the Implementation of Section 177 of the Compact and for continued waterborne transportation of agricultural products to Enewetak including operations and maintenance of the vessel used for such purposes.

Post, p. 1812.

(3) **PAYMENTS.**—Payments under this subsection shall be provided to such extent or in such amounts as are necessary for services and other assistance provided pursuant to this subsection. It is the sense of Congress that after the periods of time specified in paragraphs (1) and (2) of this subsection, consideration will be given to such additional funding for these programs as may be necessary.

(i) **RONGELAP.**—(1) Because Rongelap was directly affected by fallout from a 1954 United States thermonuclear test and because the Rongelap people remain unconvinced that it is safe to continue to live on Rongelap Island, it is the intent of Congress to take such steps (if any) as may be necessary to overcome the effects of such fallout on the habitability of Rongelap Island, and to restore Rongelap Island, if necessary, so that it can be safely inhabited. Accordingly, it is the expectation of the Congress that the Government of the Marshall Islands shall use such portion of the funds specified in Article II, section 1(e) of the subsidiary agreement for the implementation of section 177 of the Compact as are necessary for the purpose of contracting with a qualified scientist or group of scientists to review the data collected by the Department of Energy relating to radiation levels and other conditions on Rongelap Island resulting from the thermonuclear test. It is the expectation of the Congress that the Government of the Marshall Islands, after consultation with the people of Rongelap, shall select the party to review such data, and shall contract for such review and for submission of a report to the President of the United States and the Congress as to the results thereof.

Hazardous  
materials.  
Contracts.

Post, p. 1812.

Report.

(2) The purpose of the review referred to in paragraph (1) of this subsection shall be to establish whether the data cited in support of the conclusions as to the habitability of Rongelap Island, as set forth in the Department of Energy report entitled: "The Meaning of Radiation for Those Atolls in the Northern Part of the Marshall Islands That Were Surveyed in 1978", dated November 1982, are

Report.

adequate and whether such conclusions are fully supported by the data. If the party reviewing the data concludes that such conclusions as to habitability are fully supported by adequate data, the report to the President of the United States and the Congress shall so state. If the party reviewing the data concludes that the data are inadequate to support such conclusions as to habitability or that such conclusions as to habitability are not fully supported by the data, the Government of the Marshall Islands shall contract with an appropriate scientist or group of scientists to undertake a complete survey of radiation and other effects of the nuclear testing program relating to the habitability of Rongelap Island. Such sums as are necessary for such survey and report concerning the results thereof and as to steps needed to restore the habitability of Rongelap Island are authorized to be made available to the Government of the Marshall Islands.

(3) It is the intent of Congress that such steps (if any) as are necessary to restore the habitability of Rongelap Island and return the Rongelap people to their homeland will be taken by the United States in consultation with the Government of the Marshall Islands and, in accordance with its authority under the Constitution of the Marshall Islands, the Rongelap local government council.

Hazardous  
materials.

*Ante*, p. 1781.

91 Stat. 1159.  
94 Stat. 84.

(j) **FOUR ATOLL HEALTH CARE PROGRAM.**—(1) Services provided by the United States Public Health Service or any other United States agency pursuant to section 1(a) of Article II of the Agreement for the Implementation of Section 177 of the Compact (hereafter in this subsection referred to as the "Section 177 Agreement") shall be only for services to the people of the Atolls of Bikini, Eniwetok, Rongelap, and Utrik who were affected by the consequences of the United States nuclear testing program, pursuant to the program described in Public Law 95-134 and Public Law 96-205 and their descendants (and any other persons identified as having been so affected if such identification occurs in the manner described in such public laws). Nothing in this subsection shall be construed as prejudicial to the views or policies of the Government of the Marshall Islands as to the persons affected by the consequences of the United States nuclear testing program.

(2) At the end of the first year after the effective date of the Compact and at the end of each year thereafter, the providing agency or agencies shall return to the Government of the Marshall Islands any unexpended funds to be returned to the Fund Manager (as described in Article I of the Section 177 Agreement) to be covered into the Fund to be available for future use.

(3) The Fund Manager shall retain the funds returned by the Government of the Marshall Islands pursuant to paragraph (2) of this subsection, shall invest and manage such funds, and at the end of 15 years after the effective date of the Compact, shall make from the total amount so retained and the proceeds thereof annual disbursements sufficient to continue to make payments for the provision of health services as specified in paragraph (1) of this subsection to such extent as may be provided in contracts between the Government of the Marshall Islands and appropriate United States providers of such health services.

Hazardous  
materials.

*Post*, p. 1812.

(k) **ENJERI COMMUNITY TRUST FUND.**—Notwithstanding any other provision of law, the Secretary of the Treasury shall establish on the books of the Treasury of the United States a fund having the status specified in Article V of the subsidiary agreement for the implementation of Section 177 of the Compact, to be known as the



N-3        The following comments relate to the timing of the evacuation of the Rongelap people.

(a) According to C. L. Dunham, Director of the AEC Division of Biology and Medicine, (Cronkite et al, 1956), "unexpected changes in the wind structure deposited radioactive materials on inhabited atolls and on ships of Joint Task Force 7, which was conducting the tests. Radiation surveys of the areas revealed radiation levels above permissible levels: therefore evacuation was ordered, and was carried out as quickly as possible with the facilities available to the Joint Task Force".

(b) According to Merrill Eisenbud (personal communication, see references) a scientific member of the Task Force, "There are many unanswered questions about the circumstances of the 1954 fallout. It is strange that no formal investigation was ever conducted. There have been reports that the device was exploded despite an adverse meteorological forecast. It has not been explained why an evacuation capability was not standing by, as had been recommended, or why there was not immediate action to evaluate the matter when the Task Force learned (seven hours after the explosion) that the AEC Health & Safety Laboratory recording instrument on Rongerik was off scale. There was also an unexplained interval of many days before the fallout was announced to the public".

(c) Since the Rongelapese had been evacuated prior to previous tests, it is not clear why the usual procedure was changed. In February 1954, Dr. Bertell has told me, Magistrate John Angain of Rongelap was told about the Bravo test, but was not given the date. He said that "there are no orders from Washington to evacuate the people".

(d) Rongelap was evacuated on March 3, 1954, approximately 50-55 hours after the shot.

N-4

Part A of this Note deals with thyroid dosage relating to the Bravo event in 1954. It comprises two tables.

Part B consists of a letter from Dr. W. H. Adams of Brookhaven National Laboratory to Dr. Roger Ray of DOE. It deals with the question of whether or not prolonged residence on Rongelap since 1957 has resulted in an increase in thyroid neoplasia. It also considers changes in longevity and blood counts.

TABLE N.4A #1 THYROID DOSE FROM INDIVIDUAL RADIONUCLIDES  
IN FALLOUT TO THE ADULT MALE <sup>a</sup><sub>b</sub>

Source	Half-life	Per cent physical decay in 3 weeks	Dose rads
<u>Internal exposure</u>			
Iodine-135	6.6 h	100%	190 rad
Iodine-134	53.2 min	100%	3
Iodine-133	21 h	100%	550
Iodine-132	2.3 h	100%	7
Iodine-131	8.04 d	84%	130
Tellurium-131	30 h + 8.04 d	79%	120
Tellurium-131m	25 min + 8.04 d	84%	13
<u>External exposure</u>			190
<u>Total dose</u>			1203

<sup>a</sup>/ Lessard et al, (1985)

<sup>b</sup>/ Exposure to the fallout on Rongelap Island occurred for about 45 hours. The fallout fell for about 7 hours.

TABLE N.4A #2 Total Thyroid Absorbed-Dose Estimate (1954)

Age	Average Estimate, rad									
	Rongelap Island			Sifo Island			Utrik Island			
	Internal	External	Total	Internal	External	Total	Internal	External	Total	Total
Adult Male	1000	190	1200	280	110	400	150	11	160	160
Adult Female	1100	190	1300	290	110	410	160	11	170	170
Fourteen-Year-Old	1400	190	1600	410	110	530	220	11	230	230
Twelve-Year-Old	1600	190	1800	450	110	570	240	11	250	250
Nine-Year-Old	2000	190	2200	540	110	660	300	11	310	310
Six-Year-Old	2400	190	2600	640	110	760	340	11	350	350
One-Year-Old	5000	190	5200	1300	110	1400	670	11	680	680
Newborn	250	190	440	-	-	-	48	11	59	59
In Utero, 3rd tri.	680	190	870	-	-	-	98	11	110	110
In Utero, 2nd tri.	-	-	-	490	110	610	260	11	270	270
Maximum Estimate, rad										
Adult Male	4000	190	4200	1120	110	1200	600	11	610	610
Adult Female	4400	190	4600	1160	110	1300	640	11	650	650
Fourteen-Year-Old	5600	190	5800	1600	110	1700	880	11	890	890
Twelve-Year-Old	6400	190	6600	1800	110	1900	960	11	970	970
Nine-Year-Old	8000	190	8200	2200	110	2300	1200	11	1200	1200
Six-Year-Old	9600	190	9800	2600	110	2700	1400	11	1400	1400
One-Year-Old	20000	190	20000	5200	110	5300	2700	11	2700	2700
Newborn	1000	190	1200	-	-	-	190	11	200	200
In Utero, 3rd tri.	2700	190	2900	-	-	-	390	11	400	400
In Utero, 2nd tri.	-	-	-	2000	110	2100	1000	11	1000	1000

\*Multiply by 0.01 to obtain Gy.

Source: Lessard et al, 1985, p.61

N-4B

The following letter is from Dr. W. H. Adams of Brookhaven National  
National Laboratory to Dr. Roger Ray of DOE.

2108

July 18, 1985

Mr. Roger Ray  
Deputy for Pacific Operations  
Nevada Operations Office  
Department of Energy  
P.O. Box 14100  
Las Vegas, NV 89114

Dear Roger:

In view of the recent evacuation of Rongelap, which appears to have been precipitated by concern about harmful residual radioactivity on the atoll, we have reviewed our medical records to see if there is any clinical evidence that supports this conclusion and course of action.

Since 1957 an unexposed population of Marshallese of Rongelap ancestry has been examined periodically by the Brookhaven medical team. This population (the Comparison group) is similar in age and sex distribution to the exposed people of Rongelap. The reason for examination of the unexposed group has been to obtain baseline incidences of diseases in the general Marshallese population as an aid in detection of previously unidentified radiation hazards which might affect the exposed group.

Collected data on the unexposed people are sufficient to assess the effect of residence on Rongelap (since 1957) on longevity, thyroid neoplasia, and blood counts. We have done a retrospective analysis of their medical records; 133 of the group are living and 54 are deceased. We have arbitrarily selected for analysis the following divisions of years of residence on Rongelap:

Short-term - <3 years (average, 1.0 years)  
Intermediate - 4 - 14 years (average, 7.5 years)  
Long-term - >15 years (average, 20.9 years)

The place of residence for a given year is defined as the place where an individual received his medical examination. Since there is considerable migration of Marshallese among the atolls, the site of examination may not always be the same as the site of residence. Overall, however, there should be a good correlation between the two.

Mr. Roger Ray  
 July 18, 1985  
 Page 2

### Effects on Longevity

There is no evidence that prolonged residence on Rongelap since 1957 has resulted in a shortening of life expectancy:

<u>Residence Category</u>	<u>Number of Deaths</u>	<u>Mean age at Death</u>
Short-term	20	61.4 years
Intermediate	27	66.6 years
Long-term	5	70.0 years
Total	52*	Average 64.9 years

\* Does not include 2 accidental deaths.

### Effects on Thyroid Neoplasia

There is no evidence that prolonged residence on Rongelap since 1957 has resulted in an increase in thyroid neoplasia. Nine unexposed persons in the Comparison group have had surgery for thyroid nodules:

<u>Residence Category</u>	<u>Number of Persons</u>	<u>Mean Age in 1985 (yr)</u>	<u>Number with Thyroid Nodules Removed</u>	<u>Number of Thyroid Cancers</u>
Short-term	58	47.1	4 (7%)	0
Intermediate	46	46.4	3 (7%)	0
Long-term	29	46.9	2 (7%)	1
Total	133		9	1

These figures apply to the 133 unexposed persons in the Comparison group who are living. All of the 9 persons who had thyroid nodules removed are still alive.

### Effects on Blood Counts (1985 data)

There is no detectable effect of residence on Rongelap on blood counts:

<u>Residence Category</u>	<u>Number Tested</u>	<u>Neutrophils/ul ±SD</u>	<u>Lymphocytes/ul ±SD</u>	<u>Platelets/ulx10<sup>3</sup> ±SD</u>
Short-term	24	4851±2089	2754±1006	279±111
Intermediate	40	3838± 992	2835± 908	292± 59
Long-term	26	4366±1551	2612± 787	262± 51

A test of equality of means showed no statistically significant differences among the three categories. Note that the number of blood tests performed (90) is less than the number of persons in the Comparison group. This is because not all were seen in the March-April, 1985, survey.

Mr. Roger Ray  
July 18, 1985  
Page 3

We have also considered thyroid nodules and current blood cell counts as they may relate to early residence on Rongelap, since a greater radiation risk would have existed during the early years after the 1954 fallout. Thirty-four persons in the Comparison group resided in Rongelap for 4-6 years commencing with the return to the atoll in 1957. Only 1 nodule, an "occult carcinoma", has occurred in this subgroup (3.0%), whereas the other 8 nodules, including the two true thyroid carcinomas, occurred in the other 99 persons in the Comparison group (8.1%). There was also no difference in blood cell counts:

Time of Residence	Number Tested (1985)	Neutrophils/ul ±SD	Lymphocytes/ul ±SD	Platelets/ulx10 <sup>3</sup> ±SD
Early	29	4032±1543	2713±836	267±7
Late	77	4349±1599	2756±951	284±10

If you wish us to examine any other parameters do not hesitate to ask.

Sincerely yours,

William H. Adams, M.D.

WHA/elr

N-5

The sequence of safety recommendations and guides has run as follows.

(a) In 1954 the National Bureau of Standards Handbook 59 presented the recommendations of the NCRP. The maximum permissible dose to the bone marrow (and hence to the entire body) was 0.3 rem per week.

(b) In January, 1957, the whole-body dose for the general population was lowered to .5 rem per year by the NCRP. This was published as an insert into Handbook 59. The AEC also published this and other recommendations in Appendix 10, p. 400 of its 22nd Semiannual Report to the Congress.

(c) In 1960, the Federal Radiation Council defined two guides for the general population. The "radiation protection guide" for the usual case of protection was .170 rem per year. The "protective action guide", to cover spills and other accidents, was .2 rem per year to the bone marrow. These regulations, now administered by EPA, are still in force.

(d) In the period 1985-87, the ICRP (1985) and the NCRP (1987) dropped their recommendations for the general population to .1 rem per year.

When the Rongelap people returned in 1957, therefore, the guide employed by the AEC was 0.5 rem per year. It is not clear to me that this guide was met, although it may have been approximately, i.e., within a factor of two. The external dose was stated to be less than 0.5 R/year, and strontium-90 was considered to be the only significant radionuclide determining the internal dose (Dunning 1957). Lessard (Note 7), by extrapolation, found the committed effective dose equivalent to be about 0.7 rem in 1957, .44 rem in 1958, and .36 rem in 1959. These estimates do not allow for the contributions of plutonium and americium.

N-6

To be rewritten.

For the nonprofessional reader, the following is an explanation of the specific radiological meaning of the terms, exposure and dose. Very simply, the medical analogy would be this. A patient takes a spoonful of heart medicine -- radiologically considered, that is his exposure.

Of the swallowed medicine, three-quarters are eliminated but one-quarter passes from the intestine into the circulation and is absorbed by the heart -- that one-quarter is the dose. It would be expressed per gram of heart tissue.

For exposure to radiation per se, the unit is the roentgen (R), measured in air. For radionuclides (atoms which spontaneously decay and emit radiation), the units are the bequerel (Bq), equal to 1 atomic disintegration per second, or the curie (Ci),  $3.7 \times 10^{10}$  disintegrations per second. The microcurie (uCi) and the picocurie (pCi) are respectively 1 millionth of a curie, and 1 millionth of a microcurie (27 pCi equal 1 Bq).

The units of dose are the rad (for any type of ionizing radiation: 100 ergs absorbed per gram of tissue); the rem (dose equivalent in biological effect to 1 rad of standard radiation). The particular point to remember about radiation dose is that it is per gram of tissue. A whole-body dose of 100 rad means that every gram (on average) received 100 rad; it does not mean that the entire body received 100 rad to be distributed throughout the tissues.

Both exposure and dose are referred to as resulting from external or internal sources. An external exposure or external dose is the result of a radiation source outside of the body, e.g., fallout contaminated soil. An internal dose would result from a source inside of the body, e.g., radioactive iodine due to the use of fallout-contaminated drinking water.

In the case of radionuclides, we shall use the term whole-body dose in the technical sense of committed effective dose equivalent. For a particular tissue, the tissue dose would be the committed dose equivalent. Such doses can be calculated for 1 year or 30 years, etc.

Dose: in rads

Dose equivalent: in rem

Effective dose equivalent refers to the whole-body dose

Committed effective dose equivalent: whole-body dose for radionuclides in the body over a period of time



N-7

The whole-body counter measures the quantity and the energy of the gamma ray photons that have been emitted by cesium-137, or other radionuclides, and that escape from the body. In principle, the machine is calibrated by measuring the escape of gamma rays from a phantom which has been loaded with the radionuclides in question. Obviously, the whole-body counter comes closest to giving a direct measurement of the body-content. The collected data obtained with it are presented in Tables N.7, #1, #2, and #3.

In the case of radionuclides that emit beta rays (strontium-90 or alpha particles (transuranics), whose range in tissue before absorption may be at most a centimeter or so down to some micrometers, another method must be used. Recourse is had to measuring the daily radionuclide excretion in the urine. The body content is then calculated from knowledge of the metabolism of the radionuclide in question. This method is not as reliable as whole-body counting. Fortunately in the present case the detection of strontium and the transuranic elements is not as important as the detection of cesium.

The dose can also be calculated from the diet. The primary obstacle here is that the diet is very difficult to ascertain accurately, and in addition more assumptions must be made regarding the metabolism of the radionuclide than would be the case above. The Livermore results are based on this method.

Conversely, knowing the daily urinary output of a radionuclide, it is possible to calculate the daily intake by ingestion. For example, based on the work of Jones et al (1985), Skrabble et al (1987) and Moss (1988), Dr. E. T. Lessard of the Brookhaven Laboratory has calculated the factors for plutonium-239 given in Table N.4 # 4. When the daily intake is multiplied by the factor, the urinary output is obtained. Conversely, when the urinary output is known, dividing it by the factor will predict the daily intake. The Jones and Moss alternatives are offered; at 20-30 years on a constant diet, they differ by a factor of 1.75. I used the Moss-based factor for the calculations used in the text, Section 4.3, because it corrects for earlier errors in the data base which Jones did not know about.

(Cont.)

## Note 7 (cont.)

The urine data supplied by Dr. Lessard were not normally distributed:

(a)	Below $30 \times 10^{-6}$ pCi/day (the method's limit)	= 19 persons
(b)	30 - 499	= 11 persons
(c)	500 - 999	= 2 persons
(d)	999 - 3400	= 3 persons

Perhaps two or more unrecognized populations were being tested. For orientation and discussion, I therefore took the median value to represent the whole group--it would be no more than  $30 \times 10^{-6}$  pCi/day. Among the causes for the wide distribution might be technical error, but also abnormal or hitherto unrecognized physiological factors which would be of major interest to define.

I would also note that the predicted daily oral intake of plutonium-239 based on the median urine is .13 picocuries/day, not much different from the dietary estimate of .23 picocuries/day. The factor of two tends to parallel the ratio of their cesium determinations. (The activity ratio plutonium-240/plutonium-239 is 0.6.)

I understand that DOE is formulating plans to look into the matter.

TABLE N.7 #1

AVERAGE RADIONUCLIDE CONTENT AND TIME SINCE  
REHABILITATION FOR RONGELAP ADULTS

	Adult Males (>15a)		Adult Females (>15a)		Adults (>15a)		Time Post Rehabilitation Days	Year
	Body Burden Bq	Number of Individuals	Body Burden Bq	Number of Individuals	Body Burden Bq	Number of Individuals		
$^{60}\text{Co}$	$1.1 \times 10^0$	(A)	$6.3 \times 10^{-1}$	(A)	$9.3 \times 10^{-1}$	(A)	0	1957
	$3.7 \times 10^2$	37	$2.9 \times 10^2$	37	$3.3 \times 10^2$	74	1370	1961
	$9.3 \times 10^1$	45	$7.4 \times 10^1$	45	$8.1 \times 10^1$	90	2831	1965
$^{65}\text{Zn}$	$1.9 \times 10^3$	4(B)	(C)	(C)	(C)	(C)	0	1957
	$2.3 \times 10^4$	17	$6.4 \times 10^3$	8	$1.8 \times 10^4$	25	244	1958
	$1.6 \times 10^4$	30	$1.4 \times 10^4$	12	$1.5 \times 10^4$	42	304	1958
	$2.3 \times 10^4$	32	$1.9 \times 10^4$	27	$2.1 \times 10^4$	59	639	1959
	$3.5 \times 10^3$	38	$3.1 \times 10^3$	23	$3.4 \times 10^3$	61	1370	1961
$^{55}\text{Fe}$	$1.6 \times 10^4$	28	$1.5 \times 10^4$	32	$1.5 \times 10^4$	60	4626	1970
$^{90}\text{Sr}$	$7.0 \times 10^0$	(A)	$5.2 \times 10^0$	(A)	$6.3 \times 10^0$	(A)	0	1957
	$1.7 \times 10^1$	11	$1.1 \times 10^1$	4	$1.4 \times 10^1$	15	304	1958
	$4.7 \times 10^1$	24	$2.9 \times 10^1$	16	$4.1 \times 10^1$	40	639	1959
	$6.3 \times 10^1$	9	$2.5 \times 10^1$	4	$5.1 \times 10^1$	13	1370	1961
	$3.0 \times 10^2$	13	$1.8 \times 10^2$	15	$2.4 \times 10^2$	28	1696	1962
	$2.1 \times 10^2$	12	$1.9 \times 10^2$	13	$1.9 \times 10^2$	25	2100	1963
	$2.1 \times 10^2$	11	$2.0 \times 10^2$	7	$2.1 \times 10^2$	18	2466	1964
	$7.7 \times 10^1$	12	$1.6 \times 10^2$	12	$1.3 \times 10^2$	24	3561	1967
	$1.5 \times 10^2$	11	$1.2 \times 10^2$	11	$1.3 \times 10^2$	22	3927	1968
	$1.6 \times 10^2$	11	$1.3 \times 10^2$	13	$1.5 \times 10^2$	24	4292	1969
	$5.5 \times 10^1$	9	$1.5 \times 10^2$	11	$1.1 \times 10^2$	20	4657	1970
	$1.4 \times 10^2$	8	$1.2 \times 10^2$	7	$1.3 \times 10^2$	15	5022	1971
	$9.6 \times 10^1$	5	$8.7 \times 10^1$	7	$9.6 \times 10^1$	12	5388	1972
	$3.2 \times 10^2$	4	$2.1 \times 10^2$	7	$2.5 \times 10^2$	13	5753	1973
	$1.7 \times 10^2$	10	$8.5 \times 10^1$	4	$1.5 \times 10^2$	14	6118	1974
	$2.5 \times 10^2$	26	(C)	(C)	(C)	(C)	7579	1978
	$3.7 \times 10^1$	25	$2.8 \times 10^1$	19	$3.3 \times 10^1$	44	8057	1979
$^{137}\text{Cs}$	$5.2 \times 10^2$	(A)	$3.1 \times 10^2$	(A)	$4.1 \times 10^2$	(A)	0	1957
	$2.9 \times 10^4$	38	$1.9 \times 10^4$	13	$2.7 \times 10^4$	51	304	1958
	$2.9 \times 10^4$	47	$1.5 \times 10^4$	49	$2.1 \times 10^4$	96	639	1959
	$3.5 \times 10^4$	37	$1.7 \times 10^4$	37	$2.5 \times 10^4$	74	1370	1961
	$3.5 \times 10^4$	44	$1.8 \times 10^4$	45	$2.5 \times 10^4$	89	2831	1965
	$1.8 \times 10^4$	22	$1.1 \times 10^4$	24	$1.4 \times 10^4$	46	6118	1974
	$1.1 \times 10^4$	30	$7.0 \times 10^3$	21	$9.3 \times 10^3$	51	7213	1977
	$6.7 \times 10^3$	19	$5.6 \times 10^3$	18	$6.3 \times 10^3$	37	8057	1979
	$6.7 \times 10^3$	36	$7.0 \times 10^3$	30	$6.7 \times 10^3$	66	8813	1981
	$1.0 \times 10^4$	29	$7.8 \times 10^3$	18	$9.4 \times 10^3$	47	9180	1982
	$8.9 \times 10^3$	23	$7.8 \times 10^3$	29	$8.3 \times 10^3$	52	9540	1983
	$3.9 \times 10^3$	43	$3.4 \times 10^3$	35	$3.7 \times 10^3$	78	9910	1984

A = Number of individuals not recorded.

B = Measured at Argonne National Laboratory.

C = No females measured.

(This table was supplied by Dr. E. T. Lessard, Brookhaven National Laboratory)

Table N-7 #2

## BROOKHAVEN DATA FOR INTERNAL DOSE &amp; EXTERNAL EXPOSURE

Rongelap Adult Committed Effective Dose Equivalent, <sup>(1)</sup>  
Average Value Committed Each Year

$\text{mrem y}^{-1}$

microR/year

Year Post  
BRAVO

Year	<sup>60</sup> Co	<sup>137</sup> Cs	<sup>65</sup> Zn	<sup>90</sup> Sr	<sup>55</sup> Fe	Average Annual External Exposure Rate
1957	19.8	199	151	4.32	10.9	290
1958	8.35	181	33.8	3.97	8.44	210
1959	3.53	164	7.56	3.64	6.51	170
1960	1.49	149	1.69	3.34	5.02	140
1961	0.63	136	0.38	3.06	3.88	120
1962	0.27	123	0.08	2.81	2.99	100
1963	0.11	112	<u>0.02 195</u>	2.58	2.31	90
1964	0.05	102		2.37	1.78	80
1965	<u>0.02 34</u>	92.4		2.17	1.38	73
1966		83.9		1.99	1.06	66
1967		76.2		1.83	0.82	61
1968		69.2		1.68	0.63	56
1969		62.9		1.54	0.49	52
1970		57.2		1.41	0.38	49
1971		51.9		1.29	0.29	46
1972		47.2		1.19	0.22	43
1973		42.9		1.09	0.17	41
1974		38.9		1.00	0.13	38
1975		35.4		0.92	0.10	36
1976		32.1		0.84	0.08	35
1977		29.2		0.77	0.06	33
1978		26.5 1911		0.71 45	0.05 476	32 1302
1979		24.1		0.65	0.04	30 millirem
1980		21.9		0.60	0.03	29
1981		19.9		0.55	0.02	28
1982		19.1		0.50	0.02	27
1983		16.4		0.46	0.01	26
1984		14.9		0.42	0.01	25
1985		13.5		0.39	<u>0.01 14</u>	24
1986		12.3		0.36		23
1987		11.2		0.33		23
1988		10.2		0.30		22
1989		9.22		0.28		21
1990		8.38		0.25		21
1991		7.61		0.23		20
1992		6.92		0.21		19
1993		6.28		0.20		19
1994		5.71		0.18		18
1995		5.19		0.16		18
1996		4.71		0.15		17
1997		4.28		0.14		17
1998		3.89		0.13		16
1999		3.53		0.12		16
2000		3.21		0.11		15
2001		2.92		0.10		15
2002		2.65		0.09		15
2003		2.41		0.08		14
2004		2.19		0.08		14
2005		1.99		0.07		14
2006		1.80		0.06		14
2007		1.64		0.06		13
2008		1.49 245		0.05 7		13 410
2009		1.35		0.05		13 millirem

<sup>1</sup> Multiply by  $10^{-5}$  to convert to Sv.

<sup>2</sup> Multiply by 0.7 to obtain rem (whole-body).

$\Sigma$  to 1978 = 2233 + 1302 = 3535

$\Sigma$  1979-2008 = 252 + 410 = 662

This table was supplied by Dr. E. T. Leppard of the Brookhaven National Laboratory.

Table N-7 #3

SUMMARY OF BROOKHAVEN RESULTS FOR INTERNAL & EXTERNAL DOSE <sup>a/</sup>

Radionuclide	1957-78	1979-08
	mrem	mrem
<u>Internal dose</u>		
cesium-137	1911	245
strontium-90	45	7
cobalt-60	34	0
iron-55	48	0
zinc-55	195	0
Total	2,233	252
<u>External dose</u>	1,302	410

<sup>a/</sup>

Based on the data in Table N-7 #2. The external exposure rates were multiplied by 0.7 to obtain the whole-body dose. The transuranics are omitted.

TABLE N.7 #4

PLUTONIUM-239: FRACTION OF ORAL DAILY INTAKE EXCRETED IN URINE <sup>a/b/</sup>

It is assumed that the daily intake is constant over the period specified.  $F_1 = .001$ .

Elapsed interval (years)	Jones (old)	Moss (new)
1	$3.62 \times 10^{-5}$	$5.42 \times 10^{-5}$
5	$6.2 \times 10^{-5}$	--
10	$8.61 \times 10^{-5}$	$1.71 \times 10^{-4}$
20	$1.31 \times 10^{-4}$	$2.3 \times 10^{-4}$
29	$1.67 \times 10^{-4}$	$2.92 \times 10^{-4}$

<sup>a/</sup> The table's data were supplied by Dr. E. T. Lessard of the Brookhaven National Laboratory. I have used the Moss factors (Moss, 1988).

<sup>b/</sup> The intake can be calculated by dividing the urinary excretion by the factors given. For example, after 20 years of intake, the daily excretion is found to be  $3 \times 10^{-5}$  picocuries. Then the intake is:  
 $(3 \times 10^{-5}) / 2.3 \times 10^{-4} = .13$  picocuries/day.

typical of the drinking stage, it was questionably more mature, and were intermediate between drinking and copra stages. It is interesting to note that the cesium-137 mean for the 7 samples was 4.3 pCi/gram-fresh, intermediate between the drinking nut (2.3 pCi/gram) and the copra nut (6.2 pCi/gram) of previous determinations (Table 4.2 #2).

For coconut juice taken from the nuts whose meat was analyzed above, the mean for 7 samples was 1.6 pCi/gram. Previous sampling averaged about 1.3 pCi/gram (Table 4.2 #2).

(Cont.)

## Note 8 (cont.)

For 7 samples of soil (0-10 cm), the mean was 10.6 pCi/gram-dry, compared to the Livermore value of 13 pCi/gram. The original 1978 value was 12 pCi/g.

Single samples were compared in other materials. The results (pCi/gram-fresh) were (Boikat-Paretzke / Livermore): breadfruit, 4.4/3.9; arrow root, 21/17; Pandanus 26/23; lime 2.3/?.

Several analyses on single samples were done for strontium-90 and plutonium-239,-240, but I have not received the matching analyses from the Livermore Laboratory.

In the case of Ailingnae Atoll, 1 set of samples was taken on each of three islands - Mogiri, Enibuk and Gereia-Knox. Their average cesium-137 values are: drinking coconut meat, .72 pCi/gram-fresh; drinking-coconut juice, .23 pCi/gram; soil (0-10 cm), 2.7 pCi/gram-dry. The meat value is about 17% of the Rongelap Island one, the juice about 14% and the soil about 25%. Two coconut crabs averaged 1.15 pCi/gram-fresh. Their plutonium content was less than .006 pCi/gram.



N-9

The external gamma-ray exposures of Table 4.1 #1 affect all of the tissues of the body. In addition, beta rays (cesium-137 and strontium-90) emanate from soil, but have only a limited range in air and very poor penetration into the body; they might affect the body's surface in those regions which are closest to or are actually touching the ground. Shoes and clothing provide complete or almost complete protection.

External beta-ray dose is considered to be unimportant on the basis of the following. For gamma rays, the Rongelap Island/Eneu Island external-dose ratio is 1.7 (Table 4.1 #1). The beta-ray dose ratio at .007 mm depth (basal cell layer, skin) should be approximately the same. Therefore, by extrapolation from the determinations at Eneu (Shingleton et al, 1987) the Rongelap basal-cell dose would be 46 mrem/y, and at 1 cm depth practically zero (ICRP 51, Table 26). Since the radiation protection guide for skin is 5 rem/y (NCRP 1987b), the skin dose is a trivial one.

N-10

Studies on intake by inhalation concentrated on plutonium-239,-240 at Bikini Island (Shinn et al 1980). In calculating the results, it was assumed that a person would be exposed to maximum dust conditions for 5 hours per day throughout life (tilling fields), an unrealistic assumption bound to give very high exposures (tilling deposits  $1.5 \times 10^{-3}$  picocuries per hour in the lungs).

To obtain the Rongelap dose, it was assumed by Robison et al (1982b) that the distribution of particle sizes and of radionuclides was practically the same on Bikini and Rongelap Islands. Therefore, the inhalation dose on Rongelap would be to that on Bikini as the transuranic specific activity of Rongelap soil (0-5 cm) was to that of Bikini Island.

Island	Specific activity in top 5 cm of soil in 1978	Inhalation 30-year dose to bone marrow c/
	pCi/g	rem
<u>Bikini</u> a/ plutonium-239,-240 americium-241	11 8.7	.033 .035
<u>Rongelap</u> b/ plutonium-239,-240 americium-241	3.2 1.0	.010 .005

a/ Robison et al (1982a, pp. 8, 12, 44, 56).

b/ Robison et al (1982b, pp. 12, 14, 47, B10, B13).

c/ The dose throughout the bone would be about 4 times as great.

The dose is greater for a growing child. Robison et al (1982a) used a factor of 2.8 to convert the adult inhalation dose to that for the age period 0-30 years (.042 rem). The dose to the adult lung is considered to be about 2.5 times that to the marrow.

Dr. Robison (personal communication, 1988) has reviewed these dose estimates according to the more recent ICRP factors. He has reduced dust consumption by a factor of 3.5, which would reduce the dose proportionally. This is still a liberal allowance for every day of life from birth to death, but in any case a much more reasonable one. The net result is a reduction in dose for plutonium by a factor of about 3, and for americium by a factor of 4.

N-11

Diet. The major uncertainty in estimating the dose is the diet - no one knows precisely what it is. Two efforts have been made to delineate it. The first by Naidu et al (1980) (BNL 51313) was based on living experiences over the years on various Northern Marshallese Atolls and clearly demonstrated the effects of living patterns on it. Rongelap fell into their B class, one in which there was a low availability of local foods (excepting fish), overpopulation, and a good supply of imported foods (supply boat comes in regularly, say, every three weeks). Naidu et al reported the quantities of food prepared, but emphasized that they did not know how much was eaten. In any event, Robison and DOE-1982 used this estimate as the maximum level of consumption for a population.

The MLSC diet was elaborated by M. Pritchard of the Micronesian Legal Services Corporation in 1979 when he visited the Enewetak people for 2.5 weeks on Utirik Atoll (Robison et al, 1982a, UCRL-83835). His diets assumed that the supply ship came regularly, making it possible for the people to eat relatively large amounts of imported foods (rice, flour, sugar, canned goods, etc.), or that the ship did not come at all. Robison selected the adult female subgroup of the population for calculation because its consumption was greatest. DOE-1982 took this calculation for the minimal level of contaminated-food consumption.

For the MLSC diet it has been found that cesium-137 accounts for about 95% of the whole-body dose and 85% of the bone marrow dose. Strontium-90 accounts for 5% and 15%, respectively, and the transuranics for less than 1% during the first 70 years. When the supply ship is on schedule, coconut accounts for 80% or so of the radionuclide intake.

In summary, then, DOE-1982 used the Naidu type B community diet for its dose calculations. When it wished to indicate a range, it used both the type B community (high) and the MLSC diet (low). The diets are given in Table N-11 #1.

An additional fact about the preparation of fish is worth noting. The skin and bones of fish may have 50-100 times the strontium-90 specific activity of the meat. Also, the contents of the intestinal tract may be high. What is the effect of all this on dosage? First, Noshkin et al (1981) found the strontium-90 specific activities of all tissues to be below 1 pCi/g. Robison et al (personal communication, 1988), have confirmed this for mullet caught off the reef of Bikini Island (contamination levels 5-10 times those at Rongelap Island). Roast mullet and stewed mullet were tested. For stew, neither the meat, nor broth, nor skin and bones exceeded .01 pCi per gram (Table N 11.# 2). The cooking was done by natives in the customary way (the intestines were discarded).

TABLE N-11 #1 DAILY FOOD CONSUMPTION -- TWO DIETS <sup>a/</sup>

Food	Community B (adult) grams/day	MLSC Diet (adult female) grams/day
Arrowroot	0	3.9
Breadfruit	36	27.2
Banana	19	0.02
Coconut		
Drinking meat	100	--
Drinking fluid	514	--
Copra	68	--
Milk	125	--
Sprouting	100	--
Coconut "fluid"	--	142
Coconut "meat"	--	63.3
Papaya	0	6.6
Pumpkin	0	1.2
Pandanus	96	9.2
Fish	194	41.5
Eggs	--	10.7
Poultry	3	--
Wild birds	9	4.2
Domestic meat	--	21.2
Pork	1.4	--
Clams	15	8.9
Crabs	--	3.1
Octopus	20	4.5
Turtle	.1	4.3
Snails	12	--
Coconut crab	1	--
Lobster	.14	--
Shellfish	--	5.1
Total	1313.64	356.92

<sup>a/</sup> Imported foods are not included in the lists. The data are from Tables 4 and 11 in Robison et al, UCRL 52835 (1982b). Imported staples include rice (especially), sugar, flour, canned meat, canned drinks, and baby foods.

TABLE N.11 #2

STRONTIUM-90 DISTRIBUTION IN MULLET; FRESH, ROASTED,  
AND AS A STEW<sup>a/</sup>

	<u>Strontium-90, pCi/g wet weight</u>		
	Roast mullet	Mullet stew	Fresh mullet <sup>b/</sup>
Muscle (meat)	9.5 E-4	--	5.2 E-4
Bones	5.4 E-2	4.2 E-2	1.8 E-2
Duplicate bones	6.0 E-2	--	--
Skin	8.0 E-2	--	2.7 E-2
Broth	--	4.5 E-4	--
Skin + meat	--	1.8 E-3	--

<sup>a/</sup> The table was supplied by Dr. W. L. Robison of the Lawrence Livermore National Laboratory.

<sup>b/</sup> From V. Noshkin et al, UCID-20754, 1986, "Concentrations of Radionuclides in Fish Collected from Bikini Atoll between 1977 and 1984".

N-12

A major weakness in the DOE-1982 dose calculations was the small number of samples on which it was often based (URCL-52858, Pt. 1). For example, in the case of Rongelap Atoll the number of vegetation samples per island were as follows: Rongelap 35, Abar 6, Borukka 4, Mellu 6, Kabelle 6, Naen 7. On Ailingnae Atoll, there were 7 on Sifo and 2 on Uwanen.

To make up for this deficiency, vegetation specific activities were at times calculated by applying a factor to the soil's specific activity. Robison has subsequently found that such a method may give erroneous results (personal communication to H. I. Kohn).

Table N.12 #1 shows some of the inconsistencies that arise when such data are tabulated. For example, pork has the same cesium specific-activity on all islands in Rongelap Atoll; the total dose on Kabella and Mellu islands is 4.4 rem (30-year), but the internal exposures are 5500 and 8000 pCi/day, respectively.

TABLE N.12 # 1 EXPOSURE AND SPECIFIC ACTIVITY COMPARED

Location	External a/ Exposure (1978)	Total 30-year <sup>b/</sup> dose (external & internal)	Cesium-137 internal c/ exposure	Cesium specific activities in 1978 (pCi/g-fresh) <sup>d/</sup>		
				Pig: muscle, Pandanus heart	copra, cake, milk	Coconut fluid drinking meat
	$\mu\text{R}/\text{hour}$	rem	pCi/day			
<u>Rongelap Atoll</u>						
Rongelap	4.5	2.5	4300	8.5	7.6	1.4
Kabelle	14.0	4.4	5500	8.5	13.5	1.4
Mellu	--	4.4	8000	8.5	4.6	.4
Naen	43	11.0	12,100	8.5	10.9	2.6
<u>Ailingnae Atoll</u> <sup>e/</sup>						
Sifo	1.4	.5	600	1.2	1.0	.16
Ucchuwanen	1.9	1.0	1700	1.2	1.8	.43

- a/ From Figure 4.2#1 (page 31, this report)
- b/ Table 17, (UCRL 52853, Part 4), BNL community B diet, whole-body dose.
- c/ Table 14, (UCRL 52853, Part 4), cesium-137
- d/ Appendix A, (UCRL 52853, Part 4)
- e/ Ailingnae Atoll is important for food collection, especially Sifo.  
Island where a Rongelap party was visiting when the Bravo shot was fired.

## Note 13

## Comment by Consultants

Dr. Bertell and Mr. Franke have sent the following comments. I suggest that after reading them the reader review Section 5 of the Report (Discussions and Recommendations).

The fact that other consultants are not quoted does not necessarily imply their general agreement with the entire report.

It is important to bear in mind that the dosage under discussion is that from continued residence on Rongelap Island from 1978 (or the present), onwards.

I will take the liberty of commenting on four technical points which Bertell and Franke bring forward.

(1) The factor to convert roentgens (measured in air) to mean whole-body tissue dose measured in rem is 0.7. I am puzzled by Dr. Bertell's remarks on this.

(2) The .025 rem annual boundary-limit for nuclear facilities in the U.S. is based on the ALARA principle, as low as reasonably achievable. It does not apply to the totally different situation at Rongelap or Bikini, according to Dr. Alan Richardson, Chief of the Environmental Protection Agency Guides and Criteria Branch.

(3) Their reference to the United Kingdom guide being set at .05 rem/year is in error. The guide states that not more than .05 rem shall come from any one nuclear facility. The overall population guide is still .1 rem in agreement with the ICRP, according to John Dunster, recently retired Director of the U. K. National Radiation Protection Board..

(4) The cesium guide for particular food imports into the U.S. is based on the assumption that plenty of uncontaminated food is available. The decision at Rongelap rests on the average level in the whole diet, under quite different circumstances. Section 5 recommends banning arrow root for the time being, which would not be a hardship.





# INSTITUTE FOR ENERGY AND ENVIRONMENTAL RESEARCH

Washington, D.C. office:

6935 Laurel Avenue  
Takoma Park, MD 20912  
U.S.A.

Phone: (301) 270-5500  
Telex: 6502975485

## DISSENTING STATEMENT TO "PRELIMINARY REPORT, RONGELAP REASSESSMENT PROJECT", APRIL 15, 1988\*

### Abstract

The data used in the 1982 DOE bilingual report regarding the habitability of Rongelap Island was not adequate. The conclusions derived from the data which was used are incorrect. As a consequence, there is the serious possibility that radiation doses might exceed allowable levels.

The DOE report failed to acknowledge the existence of plutonium concentrations in urine of Rongelap people which exceeded expected levels. The plutonium problem is still not resolved.

The DOE declared Rongelap Island to be safe unconditionally. DOE's dose assessment is based on the assumption that a large portion of the diet consists of important food. This major assumption is omitted in the 1982 DOE report.

A complete survey of radiological conditions is recommended.

### Introduction

I was nominated as a member of Dr. Kohn's consulting team by the people of Rongelap. In my opinion, the Rongelap Reassessment Project has failed to properly or fully answer the questions asked by Congress in Public Law 99-239, section 103(i). We have an obligation to the people of Rongelap to affirm the safety and habitability of the Rongelap Atoll and that has not been done. The project should not only answer scientific questions and assess whether legal limits for radiation exposure will be exceeded or not. The Rongelap people need a level of comfort in regard to the conclusions which is beyond any doubt or uncertainty. Unfortunately, Dr. Kohn's report does not meet this objective.

My focus in the following is the amount of radiation dose from residence on Rongelap.

---

\*) The complete report was not provided at the time these comments were prepared. A more complete statement will be provided upon completion of the final report.

Statement to "Preliminary Report, Rongelap Reassessment Project", April 15, 1988  
April 15, 1988  
page 2 of 4

What did the 1982 DOE report say?

"If 233 people live on Rongelap Island and eat local food only from Rongelap Island: Scientists estimate that the largest amount of radiation a person might receive in one year from radioactive atoms that came from the U.S. bomb tests is 400 millirem. (...) The highest average amount of radiation people might receive in the coming 30 years is 2500 millirem in any part of the body and 3300 millirem in just the bone marrow ." The DOE report quotes the dose limits with 500 millirem for a single year and a total of 5000 millirem over 30 years.

Which questions did Congress ask to be reviewed?

Congress authorized a scientific determination of (1) "whether the data cited in support of the conclusions as to the habitability of Rongelap Island, as set forth in the Department of Energy report (...) are adequate" and (2) "whether such conclusions are fully supported by the data."

If either of the foregoing questions is answered in the negative, Congress has already authorized a second phase of scientific research which is to encompass "a complete survey of radiation and other effects of the nuclear testing program relating to the habitability of Rongelap Island."

Was the data used by DOE adequate?

The data used in the 1982 DOE assessment was inadequate. Aside from the fact that the assessment was based on only a small number of measurements, the problem of elevated levels of plutonium in urine of Rongelap people, known since at least 1973, was not acknowledged in the 1982 DOE report. This is a serious and significant omission.

From measurements of plutonium in urine, as imperfect as they were at that time, radiation doses exceeding DOE's regulatory limits were calculated. The concern that plutonium doses in the Marshalls might be in the tens of rems were reported to DOE representatives in a meeting in March 1981. The authors of the bilingual booklet were present. Plutonium measurements were uncertain at that time, but the degree of uncertainty was not clear. Instead of explaining the situation, the DOE opted for omission of this troublesome discovery and chose to adopt the method of dose prediction with a dietary model in the 1982 report. The investigation of plutonium levels in urine of Rongelap residents still has not been completed, almost 15 years after the initial discovery. The true plutonium dose is still not known and could well be, for some members of the Rongelap population, in excess of DOE's dose limits. (I will deal with this question below).

Were the conclusions correct?

Reviewing DOE's conclusions on the basis of the data which was used I find two major discrepancies.

First, the "maximum dose" for residents of Rongelap was given by DOE with 400 millirem per year. Rather than being the "maximum dose", this dose is referenced in the supporting documents as the 95% dose, meaning that doses for 95% of the population will be lower and for 5% of the population higher than 400 millirem.

Statement to "Preliminary Report, Rongelap Reassessment Project", April 15, 1988  
 April 15, 1988  
 page 3 of 4

According to the model used by LLNL, about 6 people would be exposed to doses above the 500 millirem per year limit quoted by the DOE.

Second, the DOE declared Rongelap Island to be unconditionally safe. However, the dietary assumptions used in the dose estimates show a high degree of imported (non-radioactive) food, thus lowering the intake of local (radioactive) food. The degree of imported food in the diet is not a natural constant but depends, among other things, on the existence of the U.S. food program which is being phased out. If habitability is defined as "possibility of full usage of Rongelap Islands natural resources for food", the Island is not habitable even by DOE's dose standards. If Rongelap people would live on local food only, for whatever reason, doses would exceed DOE's dose limits.

#### What is the radiation dose?

Suppose that the amount of local food consumed is kept at the 1978/82 level. What is the radiation dose for the Rongelap people? I agree with Dr. Kohn that the direct measurement of radioactivity in the human body is the preferred method.

However, Dr. Kohn's assessment of the average dose with 1.25 rem committed dose equivalent ("whole-body dose") over 30 years represents only one possible scenario and has two major deficiencies:

- It is based on extrapolation from the 1979 average body burden of 175,000 picocuries of cesium-137. In 1982, the average body burden was 240,000 picocuries (see Fig. 4.3#1), probably due to increased uptake of local food. Taking 1982 as the baseline, the cesium-137 dose estimate would increase from 0.62 to 0.85 rem (see Table 4.5#1).
- Kohn's estimate of plutonium dose is premature and scientifically questionable. For an accurate estimate of plutonium doses from urine data, all urine data has to be interpreted (including the data on children) and the length of residence has to be taken into account. Kohn's assumption of a 20 year continuous daily intake is not substantiated by the data and leads to underestimates of body burdens. Furthermore, at interest is the average and the maximum, not just the median dose which is referenced by Kohn.

An alternative dose estimate can be derived from the estimate of plutonium doses for the Bikini population where urine data was interpreted for a subgroup of 16 individuals which had plutonium levels above the detection limit. In these 16 cases, individual residence time was accounted for, whereas this was not the case with the Rongelap urine data. According to Dr. Lessard from Brookhaven National Laboratories, the average annual committed effective dose due to plutonium-239 is estimated with 0.25 rem. Since on Rongelap, average soil concentrations are 3.4 lower than on Bikini (see Table p.83), I would extrapolate an average plutonium dose for Rongelap people with 0.075 rem annual committed effective dose due to plutonium-239. The dose from plutonium-240 and americium-241 would be about the same. The total dose due to transuranics could well be 0.15 rem annual committed effective dose or 4.5 rem over 30 years.

- My alternative dose estimate would thus be 0.85 rem (cesium-137), 4.5 rem (transuranics), 0.021 rem (strontium-90), and 0.59 rem (external), a total of ~~5.9~~ 5.9 rem. This dose would then be above the DOE limit of 5 rem in 30 years.

Statement to "Preliminary Report, Rongelap Reassessment Project", April 15, 1988  
 April 15, 1988  
 page 4 of 4

I do not claim that my estimate represents the "true" dose. Neither is this the case with Kohn's estimate. My estimate shows that the plutonium doses might be in a region where DOE dose limits are exceeded. We will not be able to establish the "true" dose without a detailed analysis of existing urine data and a further systematic monitoring.

The above dealt with the average doses. The use of averages tends to distort the implication of radiation to real human beings. There will always be members in the population which receive more than the average. Even if the average dose could be kept below the DOE limit of 0.17 rem ~~per~~ per year (5 rem in 30 years) a segment of the population could receive doses above DOE level of 0.5 rem per year.

Would other dose limits be exceeded?

Would the radioactivity levels on Rongelap be caused from operation of a nuclear facility, the exposure would be too high since it exceeds the annual dose limit [40 CFR 190] for the maximum exposed member of the public with 0.025 rem per year (0.75 rem in 30 years).

We will have some explanation to do to the people of Rongelap why the doses they would receive are legal because they come from a nuclear weapons test fallout, whereas they would be illegal if caused by the operation of a nuclear power plant.

Current dose limits are likely to be revised in the near future. The National Radiological Protection Board in Great Britain, for example, has recently lowered the allowable doses to most highly exposed members of the public from to 0.05 millirem per year. What is an allowable dose today might soon become too high.

Levels of cesium-137 in a part of coconuts, pandanus, and arrow root harvested on Rongelap Island are exceeding limits for import into the U.S. which is currently at 10,000 pCi/kilogram. If the food is declared unsafe for the American people, how do we convince the Rongelap people that it is safe?

What is needed?

First, we need to determine what the true extent of the plutonium problem is in the Rongelap population. An extensive program of urine sampling, analysis and interpretation is needed.

Second, a program should be conducted to measure radioactivity in the whole atoll and to assess radiation exposures.

Third, measures should be taken that radiation doses from residence on Rongelap Island and food gathering on other islands in the atoll be kept as low as possible. Soil decontamination should take place on Rongelap Island as well as on the Northern islands. Special measures might have been developed to reduce the uptake of plutonium.

  
 Bernd Franke

Notes

Page 10:

There is no evidence to show that the general health of the Rongelap people has improved compared to that prior to the Bravo test in 1954. There is very little in the way of written records to use for comparison. The diabetes study was not even begun until 1974. Some disease such as venereal or vitamin A deficiency increased after the Bravo shot (Conard 1975)

Page 11:

A Rongelap youth died in 1972 from myeloid leukemia. He had been exposed to the Bravo test fallout when he was 18 months old.

There may been an artificial reduction in observed thyroid cancers attributable to surgical removal of the thvroid gland.

Page 11:

The International Institute of Concern for Public Health has asked two physicians Dr. Bernard Lau and Dr. Brenda Caloyannis, to examine health of the Rongelapese in 1985 - 1988. Their findings indicate a high level of ill health especially among those who lived on Rongelap Atoll. A separate report on this will be submitted to the U.S. Congress.

Page 12(b):

This report has not researched the various dose assignments made to the thyroid gland (1957, 1964, 1985). We are not able to conclude that the original estimates were "much too low."

Page 21(e):

According to Conard 1975 (page 16), which covers adult mortality of Rongelapese exposed and unexposed between 1956 and 1974, the first 20 years after the Bravo test:

Age Group of Deceased	Exposed No. (%)	Unexposed No. (%)
Over 60 years	12 (66.7%)	23 (74.2%)
40 to 59 years	4 (22.3%)	8 (25.8%)
Under 40 years	2 (11.1%)	-
Total	18	31

Two accidental deaths in the exposed and one accidental death in the unexposed were omitted. The death of one exposed Rongelapese with reported age 107 years apparently skewed the results so that the "average age at death" to appear similar in the two groups.

Page 14:

Although the exposed group has remained the same since 1954, the "unexposed" groups has been subjected to losses to follow-up and arbitrary increases.

Page 19:

In a situation of continuously decreasing contamination, the average dose and range of doses in the first year (which would be the highest doses) are more important than the 30 year "integral dose" calculated by Livermore. Moreover, doses to infants and children have been shown to be higher than the calculated dose to the Standard Man (Miltenberger, Lessard, Steimers and Greenhouse 1980). It is not agreed that DOE calculations were appropriated for answering the question of the Rongelap people, or for that matter, of the US Congress.

Page 19:

According to the June 1983 Bioassay Mission report of Dr. Lessard to Mr. Robert Ray, the committed effective dose equivalent from plutonium alone for those who resided on Bikini may be 350 mSv (7 mSv per year). Dr. Lessard added: "It should be noted that similar results have been obtained at Rongelap and Uterik Atolls." This dose exceeds all international and national guidelines and is extremely serious.

Bertell  
April 15, 1988

79

Page 20: Paragraph 3, line 2:

(The reference Kerr, 1980 is not given in the references page 62)

The external radiation dose is primarily due to cesium 137, with 0.66 MeV gamma radiation. The conversion to rems from external gamma radiation would be:

1 R = 0.98 rem (Radiological Health  
Handbook, Jan. 1970, US Dept. H,E, and W.)

I do not accept an arbitrary reduction of all calculations of external radiation by 30% (i.e. multiplication by 0.7)

It should also be noted that the external radiation dose one metre above the ground is inappropriate for children.

Page 23:

Reduction of the estimated 30 year transuranic whole body dose from 350 mSv (35 rem) in Lessard 1983, to 0.2 mSv (0.02 rem) in Kohn 1988, requires formal scientific explanation. The Lessard 1983 findings were based on actual urine measurements, not assumed diets.

Page 28:

Dr. Bertell does not accept the 30 year dose tabulation on page 40 because of scientific flaws noted on the previous pages. This includes but is not limited to the Kohn reduction in external doses and in dose attributable to transuranics without proper scientific evidence.

Page 30 Para.2 Line 2,ff.:

The 250 urine samples have apparently already undergone laboratory analysis. There is no justification for taking a random sample to collate. This job should be properly entered in computer together with place of residence at the time of the testing. The range should be reported and the average not the median should be used. There is no justification for using a population median to calculate collective dose. It is bad statistical practice. If the lower detectable level poses a problem it could be lowered. At any rate, urine samples with below detectable amounts of plutonium could be combined and the combined sample could be counted to obtain an average to be distributed over the samples.

Page 31:

These calculations are incorrect because of the use of a median ( as noted on page 45) and the reduction of external dose estimates (as noted on page 30). Even with these changes, the dose is for adults only and needs to be increased for infants and children.

Page 32: Second paragraph line 6:

Appealing to the incorrect calculation of adult transuranic dose (using median rather than mean) to then minimize the expected dose to children is not scientifically sound.

Page 35:

I do not accept this Table because of the errors in calculating the doses, as noted on the previous pages.

Page 33:

Given the methodological problems, statistical errors, and incomplete data, the conclusion in line 1 is not warranted at this time. The reference to Rongelapese in the second paragraph is offensive.

Page 34:

Teratogenic effects (congenital malformations) would also be expected to occur. These together with mild genetic changes would be the most frequent and most observable effects for those living on Rongelap. Choice of cancer death and severe genetic defects as the only health effects of concern reflects a legalistic, first world bias. The IICPH will submit a separate report to Congress on the observed health problems of the Rongelap people by Island of residence 1985 -88. We will also report on Rongelap children born on Majiето, Rongelap, Majuro and Ebeye in the last 15 years.



Minority Report: Rosalie Bertell, Ph.D., G.N.S.H.

The Preliminary Report, Rongelap Reassessment Project, April 15, 1988, arrived in our Toronto office 12 April 1988. The deadline for receipt of comments in California was 16 April 1988, hence these comments are necessarily incomplete and will be augmented by a separate report to Congress within the next month.

It was distressing to me to learn that blood tests and urine analyses done under US Congressional funding over the last 30 years have not even been entered into computer. No averages are available, no report has been given to the Rongelap people. The question of urine analysis for plutonium and other transuranics is serious enough to raise the question of illegal exposure of the Rongelap people to ionizing radiation even under the older more lax regulations of the 1960's. Current international opinion would be stricter by a factor of 5 to 10 times. This report glossed over the problem by selecting a sample of 35 urine reports from the 250 analyzed, and then using a median number instead of an average to extrapolate to the Rongelap people's future body burden.

The Brookhaven National Laboratory blood test data for 133 Rongelapese living on the contaminated and uncontaminated Islands has now been entered into computer. There are thirteen blood parameters for 133 people for each of 30 years (1957 - 87). It was impossible to scan these 52000 pieces of information without computerization. I fail to understand why this data has never been properly processed and analyzed, since this was obviously the purpose of collecting it. I hope to have a report on this ready within the next week.

The basic question raised by the Rongelap people and the US Congress was whether or not Rongelap Atoll is a suitable place for the Rongelap people to live, to harvest food and to bring up their children. The questions have been turned into a proliferation of numbers, many of which are not scientifically sound, which are then compared with a legalistic standard for "average consumption of food by the Standard Man". The question of pregnant women and children was not addressed, that of infants was inadequately addressed, and the fact that the Rongelapese had previous serious radiation exposure making them an already damaged people subjected to further contamination was not addressed. The IICPH will submit a separate report on these questions. It will compare the health of Rongelap children born and brought up on different Atolls.